# Hook and Line Survey of Lingcod (*Ophiodon elongatus*) and Rockfish (*Sebastes sp.*) Stocks in Southern Strait of Georgia (Statistical Areas 17, 18 and 19) October 2003

D.R. Haggarty and J.R. King

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, B.C.
V9T 6N7

2004

**Canadian Technical Report of Fisheries and Aquatic Sciences 2533**

Fisheries and Oceans / Pêches et Océans

Canada

2008-04829/

**Canadian Technical Report of Fisheries and Aquatic Sciences**

Technical reports contain scientific and technical information that contributes to existing knowledge but which is not normally appropriate for primary literature. Technical reports are directed primarily toward a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries and aquatic sciences.

Technical reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report is abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Numbers 1 - 456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457 - 714 were issued as Department of the Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Technical reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out-of-stock reports will be supplied for a fee by commercial agents.

**Rapport technique canadien des sciences halieutiques et aquatiques**

Les rapports techniques contiennent des renseignements scientifiques et techniques qui constituent une contribution aux connaissances actuelles, mais que ne sont pas normalement appropriés pour la publication dans un journal scientifique. Les rapports techniques sont destinés essentiellement à un public international et ils sont distribués à cet échelon. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du ministère des Pêches et des Océans, c'est-à-dire les scences halieutiques et aquatiques.

Les rapports techniques peuvent être cités comme des publications complètes. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports techniques sont résumés dans la revue *Résumés des sciences aquatiques et halieutiques*, et ils sont classés dans l'index annual des publications scientifiques et techniques du Ministère.

Les numéros 1 à 456 de cette série ont été publiés à titre de rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457 à 714 sont parus à titre de rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715 à 924 ont été publiés à titre de rapports techniques du Service des pêches et de la mer, ministère des Pêches et de l'Environnement. Le nom actuel de la série a été établi lors de la parution du numéro 925.

Les rapports techniques sont produits à l'échelon regional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés seront fournis contre rétribution par des agents commerciaux.

Canadian Technical Report of

Fisheries and Aquatic Sciences 2533

2004

# HOOK AND LINE SURVEY OF LINGCOD (*Ophiodon elongatus)* AND ROCKFISH *(Sebastes sp.)* STOCKS IN SOUTHERN STRAIT OF GEORGIA (STATISTICAL AREAS 17, 18 AND 19) OCTOBER 2003

By

D.R. Haggarty and J.R. King

Fisheries and Oceans Canada
Science Branch, Pacific Region
Pacific Biological Station
Nanaimo, B.C.
V9T 6N7

© Her Majesty the Queen in Right of Canada, 2004
Cat No. Fs 97-6/2533E ISSN 0706-6457

Correct citation for this publication:

Haggarty, D.R., and J.R. King. 2004. Hook and line survey of Lingcod (*Ophiodon elongatus*) and Rockfish (*Sebastes sp.*) stocks in southern Strait of Georgia (statistical areas 17, 18 and 19) October 2003. Can. Tech. Rep. Fish. Aquat. Sci. 2533: 38p.

## TABLE OF CONTENTS

LIST OF TABLES ..... iv
LIST OF FIGURES ..... v
LIST OF APPENDICES ..... vi
ABSTRACT ..... vii
RÉSUMÉ ..... viii
INTRODUCTION ..... 1
METHODS ..... 2
  ANALYSIS ..... 3
RESULTS ..... 4
  CATCH RATES ..... 4
  BIOLOGICAL DATA ..... 5
DISCUSSION ..... 5
ACKNOWLEDGEMENTS ..... 7
REFERENCES ..... 8

## LIST OF TABLES

Table 1. 2003 lingcod CPUE (LCPUE) and tc l fish CPUE (TCPUE) (number of fish per hour) data by site, date and depth strata. (Shallow=0-25 m; deep=26-50m)..... 10
Table 2. Total Catch Per Unit of Effort (TCPUE) (number of fish per hour) by fishers 1-4 is shown for repeated sites. Fisher bias for each fisher was analysed with a two-tailed Wilcoxon Signed Rank Test (non-parametric paired-t test) using repeated sites as pairs. No significant differences were observed ($p>0.05$)..... 11
Table 3. Lingcod CPUE (LCPUE) (number of fish per hour) descriptive statistics by site, statistical area, depth strata, current, tide, sea state and time. Significance was tested by Kruskal-Wallis non-parametric Anova or Mann-Whitney two-sample test..... 12
Table 4. Lingcod CPUE (number of fish per hour) was compared between years with a Mann-Whitney test in Statistical Areas 18 and 19, and among years in Statistical Area 17 with a Kruskal-Wallis test. A significant difference was observed in Statistical Area 17..... 14
Table 5. Copper Rockfish CPUE (number of fish per hour) was compared among years. For the Area 18-19 comparison, depth strata was used as a covariate in a rank-transformed ANCOVA since more copper rockfish were caught in shallow depth strata in 1993. Only the shallow depth strata is compared in Area 17 due to greater catches in the shallow strata in the 1980's and since an unbalanced data set did not allow for an ANCOVA..... 144
Table 6. Quillback Rockfish CPUE (number of fish per hour) was compared among years. For the Area 18-19 comparison, depth strata was used as a covariate in a rank-transformed ANCOVA since more quillback rockfish were caught in deep depth strata in 1993. Depth strata were pooled in Area 17 since catch rates did not vary with depth..... 15
Table 7. We tested for a difference in lingcod length between males and females using an ANOVA. Female lingcod were significantly larger than males. Accordingly, sex was used as a covariate when we used an ANCOVA to test for differences in lingcod length between depth strata and among fishers. No other significant differences in length were observed..... 15
Table 8. The length of lingcod did not vary between 1993 and 2003 in Statistical Areas 18 and 19 when the data were compared using an ANCOVA with sex as a covariate. The length of male and female lingcod did not vary between 1985 and 2003 in Statistical Area 17 when tested with an ANOVA. An ANCOVA could not be used Statistical Area 17 due to unbalanced data..... 16
Table 9. Maturity classes of male and female lingcod caught October 2003. Most lingcod were immature or maturing. (See Appendix Table 4 for a description of maturity classes)..... 16

## LIST OF FIGURES

Figure 1. Locations of sites sampled Statistical Areas 17, 18 and 19 during the 2003 Hook and Line Survey for lingcod and rockfishes.....17
Figure 2. Lingcod CPUE observed at each site in Statistical Area 17 in October 2003. **A**: shallow depth strata (0-25 m); **B**: deep depth strata (26-50 m).....18
Figure 3. Lingcod CPUE observed at each site in Statistical Areas 18 and 19 in October 2003. A: shallow depth strata (0-25 m); B: deep depth strata (26-50 m). ..... 19
Figure 4a. Box and whisker plot depicting lingcod CPUE (fish per hour) in Statistical Areas 18-19 between years. ..... 20
Figure 4b. Box and whisker plot depicting lingcod CPUE (fish per hour) in Statistical Area 17 among years.....20
Figure 5a. Box and whisker plot depicting copper rockfish CPUE (fish per hour) in Statistical Areas 18-19 between years. ..... 21
Figure 5b. Box and whisker plot dipicting copper rockfish CPUE (fish per hour) in Statistical Area 17 amoung years.....21
Figure 6a. Box and whisker plot depicting quillback rockfish CPUE (fish per hour) in Statistical Areas 18-19 between years. ..... 22
Figure 6b. Box and whisker plot depicting quillback rockfish (CPUE (fish per hour) in Statistical Area 17 amoung yearsigure .....22
Figure 7. Length frequency histogram for lingcod caught in the Hook and Line Survey in Statistical Areas 17, 18 and 19, October 2003. A: male lingcod; B: female lingcod. ..... 23
Figure 8. Lingcod length to weight relationship for male and female lingcod. The equations of the line of best fit are as follows: Male: $y = 0.002\,x^{3.391}$ $R^2$=0.960; Female: $y = 0.002\,x^{3.376}$ $R^2$=0.986. ..... 24

## LIST OF APPENDICES

Appendix Table 1. Site, location, Statistical Area (SA) and environmental characteristics for each fishing set. See appendix 5 for sea state, tide, current and weather codes. Depth strata 1=0-25m; depth strata 2=26-50 m. ..... 25

Appendix Table 2. Effort data by set (depth strata 1=0-25m; depth strata 2=26-50 m) for each fisher (F1-F5) and number of fish caught by species. Species codes are as follows: LC=lingcod, CR=copper rockfish, QR=quillback rockfish, YR=yelloweye rockfish, KG=kelp greenling, BS=buffalo sculpin, RI=red Irish lord, GS=great sculpin, CA=cabezon, PS=Pacific sanddab, SS=speckled sanddab, RS=rock sole, SD=spiny dogfish..... 28

Appendix Table 3a. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for lingcod caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m..... 31

Appendix Table 3b. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for copper rockfish caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m..... 35

Appendix Table 3c. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for quillback rockfish caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m..... 36

Appendix Table 4. Description of maturity stages for male and female lingcod. ..... 37

Appendix Table 5. Codes used for sea state, tide, current and weather. ..... 38

## ABSTRACT

Haggarty, D.R., and J.R. King. 2004. Hook and line survey of Lingcod (*Ophiodon elongatus*) and Rockfish (*Sebastes sp.*) stocks in southern Strait of Georgia (statistical areas 17, 18 and 19) October 2003. Can. Tech. Rep. Fish. Aquat. Sci. 2533: 38p.

Research fishing methods using handline gear were developed in 1985 to assess near shore reef fish abundance. Several handline surveys for rockfishes and lingcod were subsequently completed between 1985 and 1993. Historical catch per unit of effort (CPUE) data is therefore available in parts of Statistical Areas 18 and 19 from 1993, and part of Statistical Area 17 from 1985, 1987 and 1988. Between October 6-31, 2003, we revisited the sites sampled in previous surveys using similar gear and methodology and compared current to historical catch rates. This survey included new sites in the Nanaimo region of Statistical Area 17 as well as in the Southern Gulf Islands (Statistical Area 18). There were no significant differences in median lingcod catch rates in Statistical Areas 18-19 between 1993 and 2003. Median lingcod catch rates in Statistical Area 17 improved from 1987-88 and were comparable to CPUE data from 1985. Dramatic declines in copper and quillback rockfish median CPUE were observed in all Statistical Areas. We found no difference in median lingcod CPUE between depth strata sampled (0-25 m and 26-50 m) or among Statistical Areas in the 2003 survey. This report details CPUE as well as the biological data collected

## RÉSUMÉ

Haggarty, D.R., and J.R. King. 2004. Hook and line survey of Lingcod (*Ophiodon elongatus*) and Rockfish (*Sebastes sp.*) stocks in southern Strait of Georgia (statistical areas 17, 18 and 19) October 2003. Can. Tech. Rep. Fish. Aquat. Sci. 2533: 38p.

Il a été élaboré en 1985 des méthodes de pêche à la ligne à main dans le but d'évaluer l'abondance de poissons de récifs à proximité du littoral. Par la suite, plusieurs relevés de sébastes et de morue-lingue ont été effectués entre 1985 et 1993 au moyen de lignes à main. C'est ainsi que l'on dispose de données historiques sur les prises par unité d'effort (PPUE) dans certaines parties des secteurs statistiques 18 et 19, pour l'année 1993, et dans une partie du secteur statistique 17, pour les années 1985, 1987 et 1988. Du 6 au 31 octobre 2003, nous avons effectué des relevés aux endroits échantillonnés précédemment, en utilisant des engins et une méthodologie similaires et nous avons comparé les taux de capture obtenus aux taux de capture passés. Ce relevé a porté sur de nouveaux points dans la région de Nanaimo du secteur statistique 17 ainsi que dans les îles du sud du golfe (secteur statistique 18). Nous n'avons relevé aucun écart significatif des taux de capture médians de morue-lingue dans les secteurs statistiques 18 et 19, entre 1993 et 2003. Les taux de capture médians de morue-lingue dans le secteur statistique 17 se sont améliorés depuis 1987-1988 et sont comparables aux PPUE de 1985. Nous avons observé des diminutions marquées des PPUE médianes de sébaste cuivré et de sébaste à dos épineux dans tous les secteurs statistiques. Nous n'avons constaté aucune différence dans les PPUE médianes de morue-lingue entre les strates de profondeur échantillonnées (0-25 m et 26-50 m) ni entre les secteurs statistiques, lors du relevé de 2003. Le présent rapport fait état des PPUE et des données biologiques recueillies.

## INTRODUCTION

Lingcod, *Ophiodon elongatus*, populations in the Strait of Georgia have been depressed for several decades (Richards and Hand 1989; King 2001). The commercial fishery was closed in 1991. The recreational fishery, prior to 2002, was subject to regulations including an eight month winter non-retention period to protect nest guarding males, non-retention of fish less than 65 cm, a one per day bag limit, and an annual catch limits of 10 lingcod per year. In 2002, the recreational fishery was closed for the retention of lingcod as an additional measure to protect this stock (King and Surry 2000).

A stock assessment framework for lingcod (*Ophiodon elongatus*) recommended development of fishery independent sources of relative abundance to monitor changes in the Strait of Georgia lingcod population (King *et al.* 2003). One recommendation was to resume the handline surveys of nearshore reef fishes conducted in 1985, 1987-88 and 1993. In 1985, 1987 and 1988 handline surveys were developed to estimate lingcod catch per unit of effort (CPUE) and conducted in the Gulf Islands Region of Statistical Area 17 (Cass and Richards 1987; Hand and Richards 1987; Hand and Richards 1989). Fishing was conducted in November and December of 1985; and December to February in 1987 and 1988. The survey area was divided into 1 min latitude by 1 min longitude blocks, and those blocks encompassing known lingcod fishing areas were identified. Ten blocks were randomly selected and used as the fishing sites for the survey. Fishing at each site was stratified by depth: 10-25 m; 26-45 m; and 46-55 m. In 1993, a handline survey for lingcod was conducted in Statistical Areas 18 and 19 during three sampling periods, June, August and October (Yamanaka and Murie 1995). Ten fishing sites were identified by fishermen as sites with frequent lingcod catches and another ten sites were randomly selected from 1 minute latitude by 1 minute longitude blocks that encompassed rocky habitat. Fishing events were stratified by depth: 0-25 m and 25-50 m. Effort was measured as the sum of fishing time for each angler.

In all surveys, lingcod CPUE decreased as depth increased (Cass and Richards 1987, Yamanaka and Murie 1995). In 1993, lingcod catch rates were highest in October, though the difference in CPUE was not significant (Yamanaka and Murie 1995).

A handline survey was conducted in October 2003 in Statistical Areas 17, 18 and 19. Sites sampled in previous surveys were sampled in 2003 and additional sites were selected in the Nanaimo region of Statistical Area 17 as well as in Statistical Areas 18 and 19 to provide even spatial coverage. Nearshore reef fishes such as lingcod, rockfishes (*Sebastes* spp.), kelp greenling (*Hexagrammos decagrammus*), cabezon (*Scorpaenichthys marmoratus*) and spiny dogfish (*Squalus acanthias*) were caught in all survey years. The 2003 lingcod survey provides an estimate of relative abundance for these species, particularly copper rockfish (*S. caurinus*) and quillback rockfish (*S. maliger*) as well.

## METHODS

The vessel used as a platform for fishing is a 6.7 m aluminium "Lifetimer" boat equipped with twin 115-horse power engines, a depth sounder and a GPS. Fishing was conducted using the handline survey methodology developed by Richards *et al.* (1985) and Richards and Cass (1985).The fishing crew consisted of four to five research personnel, with three or four people fishing at a time. We used Zebco® Rhino® rods with Rhino® RBCXL or Shakespeare ® Tidewater ® 30LCL reels, rigged with 30 lb (13.6 kg) test mono-filament line with 25 lb (11.3 kg) test leaders. Two single Mustad #92553 size 3/0 hooks with a 6-cm spacing were used on each line. The line was separated from the leader and hooks using a large (50.8 cm or 20 inch) spreader bar with a 170 g (6 oz) mooching weight. As in previous surveys, 12-cm frozen herring were used as bait. Herring were hooked through the snout and near the dorsal fin. Previous surveys used 9-kg test mono-filament with a 7-kg leader. We used stronger line than previous surveys in order to increase our chances of landing larger lingcod. Another modification of the research methodology was the use of the spreader bar. Hook size and spacing, bait type and size, and the 170 g (6 oz) mooching weight were all consistent among surveys.

Sampling sites surveyed in 1985, 1987, 1988 and 1993 were revisited (Figure 1). Historical sites were chosen using a stratified random design. All sites represented areas of presumed lingcod (age 2+) habitat. In October of 1993, 14 sites were sampled in Statistical Areas 18 and 19. Two of these sites are now Rockfish Conservation Areas (RCA) and were therefore not fished. Eleven of the fourteen sites were re-visited. Eight of ten sites in Statistical Area 17 that were sampled in the fall or winter of 1985, 1987 and 1988 were also revisited. One site, Hidden Reef, could not be located, and another, Porlier Pass, could not be fished due to strong currents.

New sampling sites were chosen in order to extend the spatial coverage of the survey. These sites were selected for their geographic location and without *a-priori* knowledge of reef fish abundance. All sites consisted of rocky reefs of appropriate depths as indicated on nautical charts. We also tried to target areas adjacent to Rockfish Conservation Areas (RCAs). Two new sites adjacent to RCAs were selected for monitoring and at least two other historical sites were in close proximity to RCAs. An additional 11 sites in Statistical Area 17 near Nanaimo and Ladysmith were also sampled. In total, we sampled 7 sites in Statistical Area 19, 10 sites in Statistical Area 18, and 19 in Statistical Area 17. Two depth strata per site were sampled (1=0-25 m, 2=26-50 m). Because lingcod CPUE decreased with depth in all previous hook and line studies, and since we wanted to reduce the bycatch of yelloweye (*S. rubberimus*) and quillback rockfish (*S. maliger*), we did not sample a third depth strata. Seventeen depth strata-site combinations were sampled on more than one occasion to assess fisher bias (improvement with time).

Similar to previous studies, fishing effort was defined as the total fishing time of all fishers. Each fisher kept track of their fishing time using a digital stop watch strapped to the butt of the rod and represented the time the bait was on or near the bottom. Fishing

time started when the fishing gear touched the bottom and stopped whenever a fish was hooked, there was a bite, the gear become fouled on the bottom, or the line was reeled in.

Each site and depth strata was fished for a total of 60 minutes. We would adjust our position within the site if no fish were caught within 10 minutes, if we felt we were no longer in appropriate lingcod habitat, or if we were no longer within the depth stratum. Variables recorded for each set included weather conditions, tide, currents, sea state and the minimum, maximum and medium depths encountered. We stopped fishing if currents were too strong for fishing to be effective.

A catch was recorded if the fish was brought to the surface and could be identified to species. Lingcod, rockfishes and kelp greenling were sampled for fork length (mm), weight (g), sex, and stage of maturity. Fin rays of lingcod and otoliths of rockfishes were collected for age estimation. Fin rays, otoliths and scales of kelp greenling were also collected to develop ageing methodology.

## ANALYSIS

Catch Per Unit of Effort (CPUE) was calculated as the number of fish per hour for total fish catches (all species together), lingcod, copper rockfish and quillback rockfish. Effort was taken as the total fishing time of all fishers.

Species CPUE were also calculated for each fisher in order to investigate bias among fishers as well as bias over time. For the sites that were sampled on more than one occasion, fisher bias was investigated using the repeated sites as pairs in a two-tailed Wilcoxon Signed Rank Test (non-parametric paired-t test). A significant result could indicate a possible bias due to changes or improvements in fishing technique. Differences among fishers were also investigated using the Kruskal Wallis test (non-parametric ANOVA). All analyses were performed using Statistix version 7.0 software.

Differences in median CPUE between sites, depth strata and conditions (tide, current, time of day, and weather) were also analysed by the Kruskal-Wallis test (non-parametric ANOVA) or Mann-Whitney test (non-parametric t-test).

CPUE were compared between years using the Kruskal-Wallis test or Mann-Whitney test. If other variables (such as depth) significantly influenced any of the data, effects of these variables were taken into consideration either by using them as a co-variate in a rank-transformed ANOVA, or by limiting data analysis to certain depth strata.

In Statistical Area 17, there were no significant differences between months in lingcod CPUE in the 1985, 1987 or 1988 data sets, so data from November through February were pooled. These surveys also included deeper depth strata. Records deeper than 55 metres were excluded from the analysis.

Male and female lingcod lengths were compared using an ANOVA. Difference in lingcod length between depth strata and among years was investigated with an ANCOVA with sex as a co-variate. We estimated the relationship between lingcod length and weight for male and female lingcod using a non-linear least squares function ($y = ax^b$).

## RESULTS

We sampled thirty-six sites from October 6-October 31, 2003. Sampling site locations are shown in Figure 1. We repeated the shallow depth strata on two separate occasions at 10 sites, and the deep strata at 7 sites (Table 1). We fished for a total of 92.7 hours over the entire survey. Fifty-one hours were spent fishing in Statistical Area 17, 20 hours in Statistical Area 18 and 22 hours in Statistical Area 19. Data for each set are reported in Appendix Table 1. Appendix Table 2 shows the catch and effort data for each set including the effort by each fisher. Length, weight, sex and stage of maturity data for lingcod, copper rockfish (*Sebastes caurinus*) and quillback rockfish (*Sebastes maliger*) are presented in Appendix Table 3.

### CATCH RATES

We calculated total catch per unit of effort (TCPUE) and lingcod catch per unit of effort (LCPUE) for each site and depth strata (Table 1). We did not find bias in the TCPUE for any fisher when we compared catch rates at sites that were re-sampled (Table 2). This suggests that catch rates were not influenced by fishing technique over time. Similarly, we compared the total and lingcod CPUEs among the four primary fishers and found no significant differences in catch rates, indicating that the individual CPUEs are comparable. Therefore, further analyses were done using the cumulative CPUEs and not individual fisher CPUEs.

Lingcod median catch rates did not differ between the two depth strata, nor did they differ among sites or Statistical Areas (Table 3, Figure 2, and Figure 3). Current, tide and sea state did not appear to affect median catch rates (Table 3); however, we did not fish in high currents. The time of day did seem to affect catch rates as a significant difference was observed with more fish being caught in the late afternoon/evening (approximately 4 pm) (Table 3). Relatively few sets (only 5) were completed at this time due to falling daylight.

Since sites in Statistical Areas 18 and 19 were sampled in 1993 and sites in Statistical Area 17 were sampled in 1985, 1987 and 1988, we could only investigate temporal changes between 1993 and 2003 in Statistical Areas 18-19, and 1985, 1987, 1988 and 2003 in Statistical Area 17. We only used October data from the 1993 survey in Statistical Areas 18-19. There was no significant difference in 2003 lingcod catch rates in Statistical Areas 18-19 as compared to the 1993 survey data (Table 4, Figure 4a). Significant differences do exist in Statistical Area 17 when the 2003 data are compared to data from the surveys in 1985, 1988 and 1987. A post-hoc comparison revealed that

median lingcod CPUE from 1985 and 2003 are significantly higher than catch rates from 1987 and 1988; however, lingcod median CPUE does not differ between 1985 and 2003 (Table 4, Figure 4b).

The total fish CPUE was significantly lower in 2003 for all Statistical Areas. This difference is mainly attributed to differences in the catch of rockfishes. Therefore, we also compared copper and quillback rockfish catch rates among years in Statistical Areas 18 and19, and 17.

Copper rockfish median CPUE varied with depth strata in 1985, 1987, 1988 and 1993, with more fish being caught in shallow stratum. When depth was accounted for by using it as a co-variate, a significant decline in copper rockfish catch rates was apparent in both Statistical Areas (Table 5, Figure 5a and b). Likewise, we found a significant decline in quillback median catch rates in both Statistical Areas over the years (Table 6, Figure 6a and b). Quillback median CPUE was higher in the deep depth stratum in 1993; therefore, we accounted for depth in the analysis for Statistical Areas 18 and 19 by using depth as a covariate in a rank-transformed ANCOVA. Quillback catch rates did not vary between depth strata 1 and 2 in Statistical Area 17 in any year so data from both depth strata were pooled.

## BIOLOGICAL DATA

The mean length of male and female lingcod was 53.7 cm and 63.1 cm, respectively (Figure 7). Female lingcod were significantly longer than males (Table 7). There was no relationship between the length of lingcod and the depth caught when sex was used as a covariate (Table 7). The length of lingcod caught did not vary with fisher.

The mean length of lingcod caught in 2003 was also compared to 1993 in Statistical Areas 18 and 19 (Table 8). For Statistical Area 17, only data from 1985 was compared to 2003 due the low catch of lingcod in 1987 and 1988. We did not observe a difference in length between years in any Statistical Area.

The length-weight relationship for male and female lingcod was estimated using a non-linear least squares function ($y = ax^b$). Male and female lingcod exhibit a similar length-weight relationship (Figure 8).

Of the 64 male and 29 female lingcod caught, 72% and 100%, respectively, were immature or maturing (Table 9).

## DISCUSSION

Lingcod median catch rates in Statistical Areas 18 and 19 do not differ significantly between 1993 and 2003. If lingcod CPUE does track lingcod abundance, then there appears to be no change in abundance over the last ten years in Statistical Areas 18 and

19. Lingcod research catch rates in 2003 have increased in Statistical Area 17 since the 1987 and 1988 surveys. The median 2003 lingcod catch rates are similar to the 1985 lingcod catch rates. Lingcod stocks in this Statistical Area may have increased to levels comparable to levels in the mid 1980's. Hand and Richards (1989) raised the possibility that the low catches observed in 1987 and 1988, as compared to 1985, were due to seasonal differences since the 1987 and 1988 samples were from December, January and February while 1985 catch rates were from November and December. Although the January and February CPUE data were lower, we did not find a significant difference in catch rates between months in a pair-wise comparison. Seasonal differences may, however, contribute to the low catch rates in 1987 and 1988. Our survey results indicate an improvement over time in lingcod catch rates in Statistical Area 17 but not in 18 and 19. There were, however, no differences in 2003 median lingcod catch rates among the Statistical Areas sampled.

Our catch rates are comparable to previous surveys since the same sampling methodology and fishing gear was used. Although we spent a longer time fishing at each site (a total of 60 minutes versus approximately 20 minutes); all catch rates were standardized to an hour. In addition, if we did not catch any fish within about 10 minutes of fishing, we would adjust our position within the sampling site.

Copper and quillback rockfish catch rates showed statistically significant declines between sampling years in all Statistical Areas. This result may be reflective of rockfish abundance in the Strait of Georgia; however, minor changes to fishing gear and methodology may have affected catch rates.

CPUE data can be used as an index of relative abundance and compared among locations and time. Understanding how CPUE relates to actual abundance of reef fish is, however, not straight-forward (Richards and Schnute 1986). In their comparison of CPUE, as determined by research angling and visual counts from a submersible, Richards and Schnute (1986) found the relationship between CPUE and density was most reliable for the most abundant species. Further investigation concerning how CPUE relates to the density of lingcod and rockfishes should be undertaken in order to explore the relationship between CPUE and density under various abundance levels of the different reef fish species. CPUE at some reference sites could be compared to visual estimates from SCUBA surveys or towed video camera work

Other factors may influence catch rates. Environmental conditions such as tide and weather did not influence our data significantly. Time of day did affect our lingcod catch rates. Lingcod are described as "voracious predators" (Cass *et al.* 1990). Many predatory fishes have been observed to increase feeding rates during low light periods at dawn and dusk (Helfman *et al.* 1997). Lingcod may also display this behaviour. Anecdotal information suggests that moon phase may also affect catch rates. We did not, however, test this factor due to the lack of replication of moon phases.

Bait loss and competition for bait have been shown to affect CPUE in the longline fishery for halibut (*Hippoglossus stenolepis*) (Skud and Hamley 1978). We found that bait loss

was higher at some sites than at others and fishers had to check their bait often if bait loss rates were high. We don't feel as though competition for bait among lingcod and other fishes should be high since lingcod are know to be aggressive predators. Competition may be a consideration for other species particularly when other fishes are at lower densities. These factors should, however, be consistent among surveys.

## ACKNOWLEDGEMENTS

Thanks to K. Mathias, V. Hodes and P. Winchell for their hard work and to C. Fort for the use of the Lifetimer. Thank you to L. Yamanaka and J. Lochead for reviewing this work.

## REFERENCES

Cass, A.J., R.J. Beamish and G.A. McFarlane. 1990. Lingcod (*Ophiodon elongatus*). Can. Spec. Pub. Fish. Aquat. Sci. 109: 40 p.

Cass, A.J. and L.J. Richards. 1987. Lingcod Hook and Line Survey in the Gulf Island Region (Statistical Area 17) of the Strait of Georgia, October – December 1985. Can. Man. Rep. Fish. Aquat. Sci. 1934: 65 p.

Hand, C.M. and L.J. Richards. 1987. Hook and Line Survey of Lingcod (*Ophiodon elongatus*) and Rockfish (*Sebastes spp.*) Stocks in the Gulf Islands Region (MSA 17) of the Strait of Georgia, January and February, 1987. Can. Man. Rep. Fish. Aquat. Sci. 1957: 57 p.

Hand, C.M. and L.J. Richards. 1989. Hook and Line Survey of Lingcod (*Ophiodon elongatus*) Stocks in the Gulf Islands Region (MSA 17) of the Strait of Georgia, 1985 to 1988. Can. Man. Rep. Fish. Aquat. Sci. 2043: 45 p.

Helfman, G.S., B.B. Collette and D.E. Facey. 1997. The Diversity of Fishes. Blackwell Science. Toronto.

King, J.R. 2001. Assessment of Lingcod in the Strait of Georgia. Can. Sci. Ad. Sec. Res. Doc. 2001/132: 41 p.

King, J.R. and A.M. Surry. 2000. Lingcod Stock Assessment and Recommended Yield Options for 2001. Can. St. Assess. Sec. Res. Doc. 2000/164: 50 p.

King, J.R., McFarlane, G.A., and Surry, A.M. 2003. Stock assessment framework for Strait of Georgia lingcod. Can. Sci. Ad. Sec. Res. Doc., 2003/04.

Richards, L.J. and A.J. Cass. 1985. 1985 Research Catch and Effort Data on Near shore Reef-fishes in the Strait of Georgia B.C. (Statistical Areas 15 and 16). Can. Man. Rep. Fish. Aquat. Sci. 1851: 68 p.

Richards, L.J., A.J. Cass, J.R. Selsby, R. van den Broek and S.D. Loynachan. 1985. A Summary of Research Catch and Effort Data for Near shore Reef-fishes Collected during 1984 in MSA 15 and 16 of the Strait of Georgia. Can. Man. Rep. Fish. Aquat. Sci. 1833: 69 p.

Richards, L.J. and C.M. Hand. 1989. Lingcod. pp. 35-61. *In* Groundfish Stock Assessment for the West Coast of Canada in 1988 and Recommended Yield Options for 1989. J. Fargo and A.V. Tyler [eds.]. Can. Tech. Rep. Fish. Aquat. Sci. 1646: 294 p.

Richards, L.J. and J.T. Schnute. 1986. An experimental and statistical approach to the question: is CPUE an index of abundance? Can. J. Fish. Aquat. Sci. 43: 1214-1227.

Skud, B.E. and J.M. Hamley. 1978. Factors affecting longline catch and effort. International Pacific Halibut Commission Scientific Report. 64.

Yamanaka, K.L. and D.J. Murie. 1995. Hook and line surveys. pp. 69-76. *In* Groundfish Stock Assessments for the West Coast of Canada in 1994 and Recommended Yield Options for 1995. M. Stocker and J. Fargo [eds.]. Can. Tech. Rep. Fish. Aquat. Sci. 2069: 440 p.

**Table 1. 2003 lingcod CPUE (LCPUE) and total fish CPUE (TCPUE) (number of fish per hour) data by site, date and depth strata. (Shallow=0-25 m; deep=26-50m).**

| Site | Date | LCPUE Shallow | LCPUE Deep | TCPUE Shallow | TCPUE Deep |
|---|---|---|---|---|---|
| Forest Island | 06-Oct | 0 | 0 | 2.94 | 2.88 |
| Little Group | 06-Oct | 0 | 0 | 5.82 | 14.46 |
| Mouat Point | 07-Oct | 5.52 | | 7.38 | |
| Tilly Point | 07-Oct | 0 | 1.08 | 0 | 2.1 |
| Wallace Point | 07-Oct | 3.78 | 0 | 8.52 | 0.96 |
| D'arcy Island | 09-Oct | 0 | 0.9 | 0.96 | 0.9 |
| Gooch Island | 09-Oct | 8.58 | 0.84 | 9.54 | 4.98 |
| Halibut Island | 09-Oct | 0 | 0 | 0 | 1.14 |
| Isabella Island | 10-Oct | 1.98 | | 1.98 | |
| Moresby Island | 10-Oct | 1.02 | 0 | 1.02 | 0 |
| Patey Rock | 10-Oct | 0.9 | 0 | 0.9 | 2.76 |
| Russell Island | 10-Oct | 0.84 | 0.9 | 1.74 | 0.9 |
| Fane Island | 13-Oct | 0.96 | 0 | 2.82 | 0 |
| Georgson Island | 13-Oct | 1.02 | 0.96 | 3 | 5.82 |
| Taylor Point | 13-Oct | 0 | 0.96 | 0 | 3.78 |
| Tilly Point | 13-Oct | 1.8 | 2.1 | 2.7 | 2.1 |
| Imrie Island | 14-Oct | 0 | 0 | 0 | 1.98 |
| Moresby Island | 14-Oct | 0 | | 5.94 | |
| Prevost Island | 14-Oct | | 0 | | 0 |
| Forest Island | 15-Oct | 0 | 0 | 0 | 1.86 |
| Forest Island | 17-Oct | 0 | 5.4 | 0 | 7.2 |
| Imrie Island | 17-Oct | 0 | 0 | 0.96 | 2.7 |
| Little Group | 17-Oct | 0 | 2.82 | 0 | 3.72 |
| Alarm Rock | 21-Oct | 0 | 3.96 | 0 | 5.94 |
| Dionisio Point | 21-Oct | 5.7 | 1.98 | 5.7 | 1.98 |
| NE Galino | 21-Oct | 0 | 0 | 0 | 1.62 |
| Coffin Island | 22-Oct | 1.8 | 0.96 | 3.6 | 3.78 |
| Alcala Point | 23-Oct | 4.74 | 0.96 | 4.74 | 1.98 |
| Detwiller Point | 23-Oct | 0 | 0.96 | 0 | 2.82 |
| Thrasher Rock | 23-Oct | 0 | | 0 | |
| Danger Reef | 24-Oct | | 5.82 | | 10.68 |
| Noel Bay | 23-Oct | 1.98 | 0 | 3.96 | 1.02 |
| Dionisio Point | 24-Oct | 3.9 | | 3.9 | |
| NE Galino | 24-Oct | 2.76 | | 2.76 | |
| Noel Bay | 24-Oct | | 1.86 | | 1.86 |
| Danger Reef | 24-Oct | 0 | 1.02 | 0.9 | 1.98 |
| Alarm Rock | 27-Oct | 1.8 | 0 | 2.76 | 0 |
| Alcala Point | 27-Oct | 0 | | 0 | |
| Danger Reef | 27-Oct | 0 | 0.9 | 0 | 0.9 |
| Round Rock | 27-Oct | 0 | 1.02 | 0 | 1.02 |
| Entrance Island | 28-Oct | 0 | 7.68 | 0 | 7.68 |
| Malaspina Point | 28-Oct | 2.1 | | 3.18 | |
| Snake Island | 28-Oct | 0 | 0 | 0 | 0 |
| DeCourcy Island | 29-Oct | 0 | 0 | 0.96 | 0 |
| Round Rock | 29-Oct | 0 | 0 | 1.8 | 0 |
| Five Fingers | 30-Oct | 0 | 0 | 0.9 | 0.9 |
| Hudson Rocks | 30-Oct | 3.78 | 0.96 | 16.02 | 3.78 |
| Neck Point | 30-Oct | 1.92 | 0 | 3.84 | 0 |
| Douglas Island | 31-Oct | 0 | | 0 | |
| Entrance Island | 31-Oct | 0 | 0 | 4.86 | 0 |
| Grey Rock | 31-Oct | | 1.98 | | 1.98 |
| Malaspina Point | 31-Oct | | 0 | | 0 |
| Snake Island | 31-Oct | | | | |

**Table 2. Total Catch Per Unit of Effort (TCPUE) (number of fish per hour) by fishers 1-4 is shown for repeated sites. Fisher bias for each fisher was analysed with a two-tailed Wilcoxon Signed Rank Test (non-parametric paired-t test) using repeated sites as pairs. No significant differences were observed (p>0.05).**

| Set | Date | Stat Area | Site name | Depth strata | TCPUE 1 | TCPUE 2 | TCPUE 3 | TCPUE 4 |
|---|---|---|---|---|---|---|---|---|
| 43 | 21-Oct | 17 | Alarm Rock | shallow | 0 | 0 | 0 | 0 |
| 65 | 27-Oct | 17 | Alarm Rock | shallow | 0 | 4.38 | 0 | 0 |
| 51 | 23-Oct | 17 | Alcala Point | shallow | 3.72 | 5.04 | 2.64 | 0 |
| 68 | 27-Oct | 17 | Alcala Point | shallow | 3.96 | 0 | 0 | 8.46 |
| 62 | 24-Oct | 17 | Danger Reef | shallow | 0 | 0 | 0 | 0 |
| 66 | 27-Oct | 17 | Danger Reef | shallow | 0 | 0 | 0 | 0 |
| 47 | 21-Oct | 17 | Dionisio Point | shallow | 0 | 0 | 0 | |
| 58 | 24-Oct | 17 | Dionisio Point | shallow | 3.42 | 5.52 | 6.24 | 0 |
| 71 | 28-Oct | 17 | Entrance Island | shallow | 0 | 0 | 0 | |
| 88 | 31-Oct | 17 | Entrance Island | shallow | 0 | 0 | 0 | |
| 1 | 06-Oct | 19 | Forest Island | shallow | 0 | 0 | 2.76 | 0 |
| 39 | 17-Oct | 19 | Forest Island | shallow | 0 | 0 | 0 | 0 |
| 30 | 14-Oct | 19 | Imrie Island | shallow | 0 | 6.84 | 3.42 | 0 |
| 42 | 17-Oct | 19 | Imrie Island | shallow | 0 | 0 | 0 | 0 |
| 3 | 06-Oct | 19 | Little Group | shallow | 3.18 | 9.84 | 2.58 | 0 |
| 37 | 17-Oct | 19 | Little Group | shallow | 0 | 0 | 0 | 3.06 |
| 45 | 21-Oct | 17 | NE Galino | shallow | 3.66 | 3.18 | 8.82 | 8.4 |
| 60 | 24-Oct | 17 | NE Galino | shallow | 0 | 11.4 | 3.48 | 4.26 |
| 69 | 27-Oct | 17 | Round Rock | shallow | 0 | 0 | 0 | 0 |
| 79 | 29-Oct | 17 | Round Rock | shallow | 0 | 7.68 | 0 | 0 |
| 44 | 21-Oct | 17 | Alarm Rock | deep | 11.94 | 8.82 | 0 | 0 |
| 64 | 27-Oct | 17 | Alarm Rock | deep | 0 | 4.5 | 0 | 0 |
| 52 | 23-Oct | 17 | Alcala Point | deep | 3.12 | 2.88 | 3 | 22.38 |
| 67 | 27-Oct | 17 | Alcala Point | deep | 0 | 0 | 0 | 0 |
| 48 | 21-Oct | 17 | Dionisio Point | deep | 5.16 | 14.88 | 2.34 | |
| 59 | 24-Oct | 17 | Dionisio Point | deep | 0 | 4.14 | 3 | 0 |
| 2 | 06-Oct | 19 | Forest Island | deep | 0 | | 0 | 2.52 |
| 40 | 17-Oct | 19 | Forest Island | deep | 3.54 | 4.92 | 0 | 0 |
| 31 | 14-Oct | 19 | Imrie Island | deep | 3.54 | 4.02 | 0 | 0 |
| 41 | 17-Oct | 19 | Imrie Island | deep | 4.8 | 13.32 | 3.3 | 12.36 |
| 4 | 06-Oct | 19 | Little Group | deep | 8.76 | 0 | 3.18 | 0 |
| 38 | 17-Oct | 19 | Little Group | deep | 4.14 | 0 | 3 | 3 |
| 70 | 27-Oct | 17 | Round Rock | deep | 0 | 3.78 | 0 | 0 |
| 78 | 29-Oct | 17 | Round Rock | deep | 0 | | 0 | 0 |
| | | | | **Significance:** | **p=0.185** | **p=0.965** | **p=0.415** | **p=0.800** |

**Table 3. Lingcod CPUE (LCPUE) (number of fish per hour) descriptive statistics by site, statistical area, depth strata, current, tide, sea state and time. Significance was tested by Kruskal-Wallis non-parametric Anova or Mann-Whitney two-sample test.**

| | N | Mean | SD | CV | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| **Depth Strata** | | | | | | | |
| Shallow | 46 | 1.2 | 1.9 | 157 | 0.0 | 0.0 | 8.6 |
| Deep | 43 | 0.9 | 1.4 | 150 | 0.9 | 0.0 | 5.8 |
| Significance | p=0.982, U=0.0005, df=1 | | | | | | |
| **Site** | | | | | | | |
| Alarm Rock | 4 | 0.7 | 1.4 | 200 | 0.0 | 0.0 | 2.8 |
| Alcala Point | 4 | 1.1 | 0.9 | 76 | 1.4 | 0.0 | 1.8 |
| Coffin I. | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| D'arcy I. | 2 | 1.9 | 2.7 | 141 | 1.9 | 0.0 | 3.8 |
| Danger Reef | 3 | 0.6 | 1.1 | 173 | 0.0 | 0.0 | 1.9 |
| DeCourcy I. | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Detwiller Point | 2 | 2.8 | 2.6 | 93 | 2.8 | 1.0 | 4.7 |
| Dionisio Point | 4 | 1.7 | 1.7 | 97 | 3.9 | 0.0 | 3.9 |
| Douglas I. | 2 | 1.0 | 1.4 | 141 | 1.0 | 0.0 | 1.9 |
| Entrance I. | 3 | 0.3 | 0.6 | 173 | 0.0 | 0.0 | 1.0 |
| Fane I. | 2 | 0.9 | 0.01 | 2 | 0.9 | 0.9 | 0.9 |
| Five Fingers | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Forest I. | 5 | 0.2 | 0.4 | 224 | 0.0 | 0.0 | 1.0 |
| Georgeson I. | 2 | 0.5 | 0.7 | 141 | 0.5 | 0.0 | 0.9 |
| Gooch I. | 2 | 0.4 | 0.6 | 141 | 0.4 | 0.0 | 0.9 |
| Grey Rock | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Halibut I. | 2 | 4.7 | 5.5 | 117 | 4.7 | 0.8 | 8.6 |
| Hudson Rock | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Imrie I. | 4 | 2.3 | 2.3 | 97 | 1.9 | 0.0 | 5.4 |
| Isabella I. | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Little Group | 4 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Malaspina Pt. | 2 | 1.0 | 1.3 | 141 | 1.0 | 0.0 | 2.0 |
| Moresby I. | 2 | 1.0 | 1.4 | 141 | 1.0 | 0.0 | 2.0 |
| Mouat Point | 2 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| NE Galiano | 3 | 3.9 | 1.9 | 49 | 3.9 | 2.0 | 5.7 |
| Neck Point | 2 | 2.4 | 2.0 | 85 | 2.4 | 1.0 | 3.8 |
| Noel Bay | 2 | 4.3 | 2.1 | 50 | 4.3 | 2.8 | 5.8 |
| Patey Rock | 2 | 0.5 | 0.7 | 141 | 0.5 | 0.0 | 1.0 |
| Prevost I. | 1 | 0.0 | 0.0 | - | 0.0 | 0.0 | 0.0 |
| Round Rock | 4 | 0.2 | 0.5 | 200 | 0.0 | 0.0 | 0.9 |
| Russel I. | 2 | 0.4 | 0.6 | 141 | 0.4 | 0.0 | 0.9 |
| Snake I. | 2 | 1.1 | 1.5 | 141 | 1.1 | 0.0 | 2.1 |
| Taylor Point | 2 | 1.0 | 0.02 | 2 | 1.0 | 1.0 | 1.0 |
| Thrasher Rock | 2 | 0.5 | 0.7 | 141 | 0.5 | 0.0 | 0.9 |
| Tilly Point | 2 | 3.2 | 3.3 | 100 | 3.2 | 0.9 | 5.5 |
| Wallace Point | 2 | 0.5 | 0.8 | 141 | 0.5 | 0.0 | 1.1 |
| Significance | p=0.099, T=46.115, df=35 | | | | | | |
| **Statistical Area** | | | | | | | |
| 19 | 21 | 1.3 | 2.2 | 177 | 0.0 | 0.0 | 8.6 |
| 18 | 19 | 0.7 | 1.3 | 170 | 0.9 | 0.0 | 5.5 |
| 17 | 49 | 1.1 | 1.6 | 141 | 0.0 | 0.0 | 5.8 |
| Significance | p=0.833, T=0.365, df=2 | | | | | | |

Table 3 (continued)

| | N | Mean | SD | CV | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| **Current** | | | | | | | |
| None | 14 | 0.5 | 0.8 | 150 | 0.0 | 0.0 | 2.0 |
| Weak | 52 | 0.9 | 1.4 | 153 | 0.0 | 0.0. | 5.8 |
| Moderate | 23 | 1.8 | 2.4 | 135 | 0.8 | 0.0 | 8.6 |
| Significance | p=0.276, T=2.576, df=2 | | | | | | |
| **Tide** | | | | | | | |
| Ebb | 23 | 0.6 | 1.1 | 196 | 0.0 | 0.0 | 3.8 |
| Flood | 46 | 1.3 | 2.0 | 146 | 0.9 | 0.0 | 8.6 |
| High Slack | 9 | 1.5 | 2.0 | 133 | 0.9 | 0.0 | 5.4 |
| Low Slack | 11 | 0.7 | 0.8 | 124 | 0.0 | 0.0 | 2.0 |
| Significance | p=0.218, T=4.438, df=3 | | | | | | |
| **Sea State** | | | | | | | |
| Calm | 40 | 1.5 | 1.8 | 126 | 0.9 | 0.0 | 5.8 |
| Ripple | 27 | 0.6 | 1.0 | 156 | 0.0 | 0.0 | 3.8 |
| Chop | 13 | 1.1 | 2.4 | 217 | 0.0 | 0.0 | 8.6 |
| Swell | 9 | 0.6 | 1.2 | 197 | 0.0 | 0.0 | 3.8 |
| Significance | p=0.134, T=5.58, df=3 | | | | | | |
| **Time of day** | | | | | | | |
| 800 | 2 | 1.0 | 1.4 | 141 | 1.0 | 0.0 | 2.0 |
| 900 | 9 | 0.9 | 1.1 | 121 | 0.0256 | 0.0 | 4.0 |
| 1000 | 11 | 0.2 | 0.6 | 332 | 0.0 | 0.0 | 2.1 |
| 1100 | 15 | 0.6 | 1.0 | 163 | 0.0 | 0.0 | 2.8 |
| 1200 | 12 | 1.0 | 2.4 | 238 | 0.0 | 0.0 | 8.6 |
| 1300 | 12 | 1.8 | 2.2 | 124 | 0.9 | 0.0 | 5.7 |
| 1400 | 12 | 1.2 | 1.4 | 112 | 0.9 | 0.0 | 4.7 |
| 1500 | 8 | 0.6 | 0.7 | 122 | 0.4 | 0.0 | 2.0 |
| 1600 | 5 | 3.6 | 2.2 | 61 | 3.9 | 0.9 | 5.8 |
| Significance | **p=0.024**, T=17.61, df=8 | | | | | | |

**Table 4. Lingcod CPUE (number of fish per hour) was compared between years with a Mann-Whitney test in Statistical Areas 18 and 19, and among years in Statistical Area 17 with a Kruskal-Wallis test. A significant difference was observed in Statistical Area 17.**

| | N | Mean | SD | CV | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| **Areas 18-19** | | | | | | | |
| 1993 | 80 | 1.9 | 3.2 | 170 | 0.0 | 0.0 | 17.9 |
| 2003 | 40 | 1.0 | 1.8 | 180 | 0.0 | 0.0 | 8.6 |
| **Significance** | p=0.783, U=0.076, df=1 | | | | | | |
| **Area 17** | | | | | | | |
| 1985 | 38 | 1.7 | 3.4 | 201 | 0.0 | 0.0 | 12.8 |
| 1987 | 104 | 0.5 | 1.5 | 327 | 0.0 | 0.0 | 6.0 |
| 1988 | 71 | 0.1 | 0.5 | 366 | 0.0 | 0.0 | 2.0 |
| 2003 | 24 | 1.8 | 1.9 | 104 | 1.4 | 0.0 | 5.8 |
| **Significance** | **p=<0.0001**, T=48.4, df=3; 2003 is significantly different from 1987 and 1988 but not 1985. | | | | | | |

**Table 5. Copper Rockfish CPUE (number of fish per hour) was compared among years. For the Area 18-19 comparison, depth strata was used as a covariate in a rank-transformed ANCOVA since more copper rockfish were caught in shallow depth strata in 1993. Only the shallow depth strata is compared in Area 17 due to greater catches in the shallow strata in the 1980's and since an unbalanced data set did not allow for an ANCOVA.**

| | N | Mean | SD | CV | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| **Areas 18-19** | | | | | | | |
| **1993** | 80 | 5.1 | 5.9 | 117 | 3.0 | 0.0 | 27.5 |
| **2003** | 40 | 0.5 | 1.2 | 220 | 0.0 | 0.0 | 6.0 |
| **Significance** | **p=<0.0001**, F=26.31, df=1, 117 | | | | | | |
| **Area 17** | | | | | | | |
| **1985** | 14 | 4.1 | 5.5 | 136 | 1.1 | 0.0 | 16.0 |
| **1987** | 45 | 6.9 | 8.1 | 118 | 5.0 | 0.0 | 33.4 |
| **1988** | 34 | 3.1 | 2.9 | 95 | 3.0 | 0.0 | 10.0 |
| **2003** | 26 | 0.5 | 0.9 | 192 | 0.0 | 0.0 | 3.9 |
| **Significance** | **p=0.0005**, T=17.881, df=3; 2003 significantly different from all other years | | | | | | |

**Table 6. Quillback Rockfish CPUE (number of fish per hour) was compared among years. For the Area 18-19 comparison, depth strata was used as a covariate in a rank-transformed ANCOVA since more quillback rockfish were caught in deep depth strata in 1993. Depth strata were pooled in Area 17 since catch rates did not vary with depth.**

| | N | Mean | SD | CV | Median | Minimum | Maximum |
|---|---|---|---|---|---|---|---|
| **Areas 18-19** | | | | | | | |
| **1993** | 80 | 6.5 | 9.6 | 149 | 3.0 | 0.0 | 62.7 |
| **2003** | 40 | 0.2 | 0.5 | 233 | 0.0 | 0.0 | 1.9 |
| **Significance** | **p=<0.0001**, F=53.28, df=1, 117 | | | | | | |
| **Area 17** | | | | | | | |
| **1985** | 42 | 6.5 | 9.8 | 152 | 3.0 | 0.0 | 46.1 |
| **1987** | 128 | 4.4 | 6.6 | 150 | 0.0 | 0.0 | 36.0 |
| **1988** | 86 | 2.7 | 3.8 | 140 | 0.9 | 0.0 | 15.0 |
| **2003** | 49 | 0.4 | 1.5 | 324 | 0.0 | 0.0 | 9.4 |
| **Significance** | **p=<0.0001**, T=22.88, df=3; 2003 is significantly different from all other years | | | | | | |

**Table 7. We tested for a difference in lingcod length between males and females using an ANOVA. Female lingcod were significantly larger than males. Accordingly, sex was used as a covariate when we used an ANCOVA to test for differences in lingcod length between depth strata and among fishers. No other significant differences in length were observed.**

| | | Male | Female |
|---|---|---|---|
| | N | 68 | 29 |
| | Mean ± SD | 536.6 ± 95.6 | 630.8 ± 116.9 |
| | Range | 368.0-830.0 | 446.0-868.0 |
| Significance | | **p=0.0001**, F=17.24, df=1, 95 | |
| **Depth Strata** | | | |
| **Shallow** | N | 44 | 13 |
| | Mean ± SD | 545.8 ± 93.7 | 600.2 ± 103.2 |
| | Range | 369.0-830.0 | 471.0-815.0 |
| **Deep** | N | 24 | 16 |
| | Mean ± SD | 519.6 ± 98.6 | 655.6 ± 124.6 |
| | Range | 368.0-760.0 | 446.0-868.0 |
| Significance | | p=0.9845, F=0.00, df=1, 94 | |
| **Fisher** | | | |
| **F1** | N | 13 | 10 |
| | Mean ± SD | 558.4 ± 132.3 | 611.3 ± 107.3 |
| | Range | 405.0-830.0 | 446.0-760.0 |
| **F2** | N | 22 | 8 |
| | Mean ± SD | 513.8 ± 82.2 | 664.1 ± 121.0 |
| | Range | 435.0-743.0 | 546.0-856.0 |
| **F3** | N | 19 | 5 |
| | Mean ± SD | 546.6 ± 93.1 | 589.2 ± 14.6 |
| | Range | 369.0-755.0 | 510.0-712.0 |
| **F4** | N | 13 | 5 |
| | Mean ± SD | 542.7 ± 82.3 | 621.0 ± 150.9 |
| | Range | 368.0-645.0 | 480.0-868.0 |
| Significance | | p=0.9059, F=0.26, df=4, 91 | |

**Table 8. The length of lingcod did not vary between 1993 and 2003 in Statistical Areas 18 and 19 when the data were compared using an ANCOVA with sex as a covariate. The length of male and female lingcod did not vary between 1985 and 2003 in Statistical Area 17 when tested with an ANOVA. An ANCOVA could not be used Statistical Area 17 due to unbalanced data.**

| Areas 18-19 | | Both Sexes | |
|---|---|---|---|
| 1993 | N | 45 | |
| | Mean ± SD | 622.8 ± 87.0 | |
| | Range | 262.0-811.0 | |
| 2003 | N | 38 | |
| | Mean ± SD | 540.1 ± 92.6 | |
| | Range | 368.0-830.0 | |
| Significance | | p=0.50, F=0.44, df=1, 79 | |
| Area 17 | | Females | Males |
| 1985 | N | 10 | 9 |
| | Mean ± SD | 572.3 ± 183.3 | 543.9 ± 93.8 |
| | Range | 352.0-932.0 | 417.0-753.0 |
| 2003 | N | 20 | 38 |
| | Mean ± SD | 642.3 ± 134.1 | 549.8 ±98.9 |
| | Range | 446.0-868.0 | 405.0-760.0 |
| Significance | | p=0.24, F=1.42, df=1, 28 | p=0.87, F=0.03, df=1, 45 |

**Table 9. Maturity classes of male and female lingcod caught October 2003. Most lingcod were immature or maturing. (See Appendix Table 4 for a description of maturity classes).**

| | Male | | Female | |
|---|---|---|---|---|
| Maturity Class | Frequency | Percent % | Frequency | Percent % |
| 1-Immature | 18 | 28.1 | 9 | 31.0 |
| 2-Maturing-small | 28 | 43.8 | 9 | 31.0 |
| 3-Maturing-large | 15 | 23.4 | 11 | 37.9 |
| 4-Mature | 2 | 3.1 | 0 | |
| 5-Ripe | 1 | 1.6 | 0 | |
| Total | 64 | 100 | 29 | 100 |




**Figure 1. Locations of sites sampled Statistical Areas 17, 18 and 19 during the 2003 Hook and Line Survey for lingcod and rockfishes.**




Figure 2. Lingcod CPUE observed at each site in Statistical Area 17 in October 2003. **A**: shallow depth strata (0-25 m); **B**: deep depth strata (26-50 m).




Figure 3. Lingcod CPUE observed at each site in Statistical Areas 18 and 19 in October 2003. **A**: shallow depth strata (0-25 m); **B**: deep depth strata (26-50 m).




**Figure 4a. Box and whisker plot depicting lingcod CPUE (fish per hour) in Statistical Areas 18-19 between years.**




**Figure 4b. Box and whisker plot depicting lingcod CPUE (fish per hour) in Statistical Area 17 among years.**




**Figure 5a. Box and whisker plot depicting copper rockfish CPUE (fish per hour) in Statistical Areas 18-19 between years.**




**Figure 5b. Box and whisker plot depicting copper rockfish CPUE (fish per hour) in Statistical Area 17 among years**




**Figure 6a. Box and whisker plot depicting quillback rockfish CPUE (fish per hour) in Statistical Areas 18-19 between years.**




**Figure 6b. Box and whisker plot depicting quillback rockfish CPUE (fish per hour) in Statistical Area 17 among years.**




**Figure 7. Length frequency histogram for lingcod caught in the Hook and Line Survey in Statistical Areas 17, 18 and 19, October 2003. A: male lingcod; B: female lingcod.**




**Figure 8. Lingcod length to weight relationship for male and female lingcod. The equations of the line of best fit are as follows: Male: $y = 0.002\,x^{3.391}$ $R^2$=0.960; Female: $y = 0.002\,x^{3.376}$ $R^2$=0.986.**

**Appendix Table 1. Site, location, Statistical Area (SA) and environmental characteristics for each fishing set. See appendix 5 for sea state, tide, current and weather codes. Depth strata 1=0-25m; depth strata 2=26-50 m.**

| Set | Site # | Site Name | SA | Depth Strata | Latitude | | | Longitude | | | Sea State | Tide | Current | Weather | Med. Depth | Start Time | End Time |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Deg | Min | Sec | Deg | Min | Sec | | | | | | | |
| 1 | 207 | Forest I. | 19 | 1 | 48 | 39 | 44.5 | 123 | 20 | 29.0 | 1 | 2 | 2 | 4 | 18 | 11:20 | 12:10 |
| 2 | 207 | Forest I. | 19 | 2 | 48 | 39 | 44.5 | 123 | 20 | 29.0 | 1 | 2 | 3 | 4 | 45 | 13:20 | 14:20 |
| 3 | 104 | Little Grp. | 19 | 1 | 48 | 40 | 10.0 | 123 | 22 | 58.1 | 2 | 2 | 1 | 2 | 15 | 14:55 | 15:36 |
| 4 | 104 | Little Grp. | 19 | 2 | 48 | 40 | 10.0 | 123 | 22 | 58.1 | 2 | 2 | 1 | 4 | 32 | 15:44 | 16:30 |
| 5 | 205 | Mouat Pt. | 18 | 1 | 48 | 46 | 43.8 | 123 | 19 | 17.5 | 1 | 4 | 2 | 3 | 13 | 9:30 | 10:22 |
| 6 | 205 | Mouat Pt. | 18 | 2 | 48 | 46 | 43.8 | 123 | 19 | 17.5 | 2 | 4 | 3 | 3 | 41 | 10:25 | 11:48 |
| 7 | 208 | Wallace Pt. | 18 | 1 | 48 | 44 | 10.1 | 123 | 14 | 4.3 | 1 | 2 | 1 | 5 | 16 | 12:22 | 12:51 |
| 8 | 208 | Wallace Pt. | 18 | 2 | 48 | 44 | 10.1 | 123 | 14 | 4.3 | 1 | 2 | 2 | 5 | 38 | 13:00 | 15:30 |
| 9 | 314 | Tilly Pt. | 18 | 1 | 48 | 43 | 57.0 | 123 | 12 | 0.4 | 1 | 2 | 2 | 1 | 14 | 16:00 | 17:00 |
| 10 | 203 | D'arcy I. | 19 | 1 | 48 | 32 | 21.3 | 123 | 17 | 13.5 | 2 | 1 | 3 | 4 | 18 | 9:20 | 10:40 |
| 11 | 203 | D'arcy I. | 19 | 2 | 48 | 32 | 21.3 | 123 | 17 | 13.5 | 3 | 4 | 1 | 4 | 30 | 10:50 | 12:00 |
| 12 | 312 | Halibut I. | 19 | 1 | 48 | 37 | 12.6 | 123 | 16 | 8.8 | 3 | 2 | 2 | 4 | 15 | 12:00 | 13:41 |
| 13 | 312 | Halibut I. | 19 | 2 | 48 | 37 | 12.6 | 123 | 16 | 8.8 | 3 | 2 | 2 | 4 | 40 | 13:49 | 14:35 |
| 14 | 110 | Gooch I. | 19 | 1 | 48 | 39 | 42.0 | 123 | 17 | 54.8 | 3 | 2 | 1 | 4 | | 14:49 | 15:38 |
| 15 | 110 | Gooch I. | 19 | 2 | 48 | 39 | 42.0 | 123 | 17 | 54.8 | 1 | 2 | 1 | 4 | 33 | 15:40 | 16:40 |
| 16 | 103 | Russell I. | 18 | 1 | 48 | 44 | 44.6 | 123 | 24 | 32.1 | 3 | 1 | 1 | 4 | 14 | 9:20 | 10:04 |
| 17 | 103 | Russell I. | 18 | 2 | 48 | 45 | 1.2 | 123 | 24 | 3.0 | 3 | 1 | 1 | 3 | 33 | 10:15 | 11:03 |
| 18 | 210 | Isabella I. | 18 | 1 | 48 | 43 | 42.0 | 123 | 25 | 50.8 | 2 | 4 | 1 | 3 | 12 | 11:18 | 12:00 |
| 19 | 210 | Isabella I. | 18 | 2 | 48 | 43 | 42.0 | 123 | 25 | 50.8 | 2 | 4 | 1 | 3 | 37 | 12:05 | 13:05 |
| 20 | 313 | Patey Rk. | 18 | 2 | 48 | 42 | 7.3 | 123 | 31 | 17.1 | 2 | 2 | 1 | 3 | 30 | 13:30 | 13:56 |
| 21 | 313 | Patey Rk. | 18 | 1 | 48 | 42 | 7.3 | 123 | 31 | 17.1 | 1 | 2 | 1 | 3 | 17 | 14:01 | 14:30 |
| 22 | 205 | Moresby I. | 19 | 1 | 48 | 43 | 57.0 | 123 | 19 | 56.5 | 2 | 2 | 1 | 3 | 15 | 15:15 | 15:53 |
| 23 | 314 | Tilly Pt. | 18 | 2 | 48 | 43 | 57.0 | 123 | 12 | 0.4 | 1 | 3 | 1 | 4 | 33 | 9:30 | 10:45 |
| 24 | 109 | Taylor Pt. | 18 | 1 | 48 | 45 | 47.0 | 123 | 7 | 41.4 | 1 | 1 | 1 | 4 | 13 | 11:00 | 11:40 |
| 25 | 109 | Taylor Pt. | 18 | 2 | 48 | 45 | 47.0 | 123 | 7 | 41.4 | 1 | 1 | 1 | 4 | 38 | 11:50 | 13:04 |
| 26 | 317 | Georgson I. | 18 | 2 | 48 | 50 | 38.9 | 123 | 14 | 1.1 | 1 | 4 | 0 | 4 | 31 | 13:35 | 14:15 |
| 27 | 317 | Georgson I. | 18 | 1 | 48 | 50 | 38.9 | 123 | 14 | 1.1 | 1 | 4 | 0 | 4 | 12 | 14:15 | 15:11 |
| 28 | 315 | Fane I. | 18 | 1 | 48 | 48 | 22.9 | 123 | 15 | 59.9 | 2 | 2 | 1 | 1 | 12 | 15:30 | 16:04 |
| 29 | 315 | Fane I. | 18 | 2 | 48 | 48 | 22.9 | 123 | 15 | 59.9 | 1 | 2 | 0 | 1 | 28 | 16:15 | 16:45 |
| 30 | 204 | Imrie I. | 19 | 1 | 48 | 41 | 36.5 | 123 | 20 | 5.8 | 3 | 2 | 1 | 1 | 9 | 9:00 | 9:42 |

| Set | Site # | Site Name | SA | Depth Strata | Latitude | | | Longitude | | | Sea State | Tide | Current | Weather | Med. Depth | Start Time | End Time |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Deg | Min | Sec | Deg | Min | Sec | | | | | | | |
| 31 | 204 | Imrie I. | 19 | 2 | 48 | 41 | 36.5 | 123 | 20 | 5.8 | 1 | 3 | 1 | 1 | 31 | 10:04 | 10:50 |
| 32 | 205 | Moresby I. | 19 | 2 | 48 | 43 | 57.0 | 123 | 19 | 56.5 | 3 | 1 | 1 | 1 | 30 | 11:05 | 11:50 |
| 33 | 316 | Prevost I. | 18 | 1 | 48 | 49 | 50.0 | 123 | 24 | 20.0 | 1 | 1 | 1 | 1 | 20 | 13:45 | 14:29 |
| 35 | 207 | Forest I. | 19 | 2 | 48 | 59 | 27.3 | 123 | 32 | 53.3 | 1 | 2 | 3 | 2 | 30 | 10:20 | 11:14 |
| 36 | 207 | Forest I. | 19 | 1 | 48 | 59 | 27.3 | 123 | 32 | 53.3 | 3 | 2 | 3 | 2 | | | |
| 37 | 104 | Little Grp. | 19 | 1 | 49 | 1 | 6.3 | 123 | 34 | 31.1 | 2 | 2 | 2 | 4 | 17 | 10:20 | 10:50 |
| 38 | 104 | Little Grp. | 19 | 2 | 49 | 1 | 3.0 | 123 | 34 | 8.0 | 2 | 2 | 1 | 4 | 31 | 11:15 | 11:40 |
| 39 | 207 | Forest I. | 19 | 1 | 48 | 59 | 8.3 | 123 | 45 | 19.3 | 1 | 2 | 1 | 4 | 19 | 11:55 | 12:23 |
| 40 | 207 | Forest I. | 19 | 2 | 48 | 59 | 2.9 | 123 | 45 | 41.2 | 1 | 3 | 2 | 2 | 40 | 12:20 | 13:00 |
| 41 | 204 | Imrie I. | 19 | 2 | 49 | 0 | 12.8 | 123 | 35 | 19.3 | 1 | 3 | 1 | 2 | 40 | 13:10 | 14:20 |
| 42 | 204 | Imrie I. | 19 | 1 | 49 | 0 | 11.1 | 123 | 35 | 21.3 | 2 | 1 | 1 | 2 | 12 | 14:28 | 14:52 |
| 43 | 53 | Alarm Rk. | 17 | 1 | 49 | 8 | 56.7 | 123 | 38 | 30.5 | 1 | 2 | 1 | 4 | 23 | 10:55 | 11:49 |
| 44 | 53 | Alarm Rk. | 17 | 2 | 49 | 9 | 15.6 | 123 | 38 | 28.6 | 1 | 2 | 1 | 4 | 35 | 11:49 | 12:35 |
| 45 | 30 | NE Galiano | 17 | 1 | 49 | 3 | 44.0 | 123 | 37 | 22.0 | 1 | 2 | 1 | 4 | 7 | 13:15 | 14:10 |
| 46 | 30 | NE Galiano | 17 | 2 | 49 | 3 | 41.7 | 123 | 37 | 6.8 | 1 | 2 | 2 | 4 | 30 | 14:15 | 15:15 |
| 47 | 27 | Dionisio Pt. | 17 | 1 | 49 | 2 | 46.3 | 123 | 36 | 14.2 | 1 | 3 | 2 | 4 | 17 | 15:30 | 16:30 |
| 48 | 27 | Dionisio Pt. | 17 | 2 | 49 | 2 | 24.0 | 123 | 35 | 42.0 | 1 | 3 | 2 | 3 | 38 | 16:35 | 17:49 |
| 49 | 301 | Coffin I. | 17 | 1 | 49 | 3 | 20.4 | 123 | 43 | 6.1 | 3 | 2 | 1 | 4 | 17 | 11:30 | 12:13 |
| 50 | 301 | Coffin I. | 17 | 2 | 49 | 3 | 26.2 | 123 | 43 | 5.8 | 3 | 2 | 1 | 4 | 39 | 12:20 | 13:20 |
| 51 | 19 | Alcala Pt. | 17 | 1 | 48 | 57 | 34.3 | 123 | 40 | 44.1 | 2 | 4 | 1 | 1 | 10 | 9:10 | 9:55 |
| 52 | 19 | Alcala Pt. | 17 | 2 | 48 | 57 | 45.3 | 123 | 40 | 51.2 | 2 | 4 | 1 | 1 | 45 | 9:55 | 11:11 |
| 53 | 3 | Thrasher Rk. | 17 | 1 | 49 | 6 | 54.9 | 123 | 47 | 47.4 | 1 | 2 | 1 | 1 | 8 | 12:12 | 12:41 |
| 54 | 3 | Thrasher Rk. | 17 | 2 | 49 | 6 | 44.3 | 123 | 47 | 24.3 | 2 | 2 | 1 | 1 | 34 | 12:50 | 13:51 |
| 55 | 12 | Detwiller Pt. | 17 | 1 | 49 | 12 | 31.0 | 123 | 48 | 58.0 | 1 | 2 | 1 | 1 | 8 | 14:15 | 15:13 |
| 56 | 12 | Detwiller Pt. | 17 | 2 | 49 | 12 | 39.2 | 123 | 48 | 14.8 | 1 | 2 | 1 | 1 | 30 | 15:20 | 15:52 |
| 57 | 13 | Noel Bay | 17 | 2 | 49 | 11 | 26.0 | 123 | 52 | 28.0 | 1 | 2 | 1 | 1 | 31 | 16:05 | 17:25 |
| 58 | 27 | Dionisio Pt. | 17 | 1 | 49 | 12 | 40.5 | 123 | 53 | 5.3 | 2 | 4 | 2 | 4 | 11 | 11:22 | 12:24 |
| 59 | 27 | Dionisio Pt. | 17 | 2 | 49 | 5 | 34.5 | 123 | 43 | 7.1 | 2 | 2 | 1 | 1 | 40 | 12:37 | 13:30 |
| 60 | 30 | NE Galiano | 17 | 1 | 49 | 5 | 33.3 | 123 | 42 | 59.7 | 1 | 2 | 1 | 1 | 8 | 13:40 | 14:20 |
| 61 | 13 | Noel Bay | 17 | 1 | 49 | 13 | 49.7 | 123 | 55 | 2.5 | 1 | 2 | 1 | 1 | 15 | 14:30 | 15:17 |
| 62 | 51 | Danger Rf. | 17 | 1 | 49 | 13 | 52.7 | 123 | 55 | 11.0 | 1 | 2 | 0 | 4 | 12 | 15:30 | 16:20 |
| 63 | 51 | Danger Rf. | 17 | 2 | 49 | 13 | 56.3 | 123 | 55 | 26.0 | 1 | 2 | 0 | 4 | 30 | 16:20 | 16:59 |
| 64 | 53 | Alarm Rk. | 17 | 2 | 49 | 13 | 39.2 | 123 | 55 | 40.6 | 1 | 1 | 1 | 4 | 39 | 9:00 | 9:40 |

| Set | Site # | Site Name | SA | Depth Strata | Latitude | | | Longitude | | | Sea State | Tide | Current | Weather | Med. Depth | Start Time | End Time |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | Deg | Min | Sec | Deg | Min | Sec | | | | | | | |
| 65 | 53 | Alarm Rk. | 17 | 1 | 49 | 14 | 15.3 | 123 | 57 | 15.3 | 1 | 1 | 1 | 4 | 14 | 9:45 | 10:30 |
| 66 | 51 | Danger Rf. | 17 | 1 | 49 | 14 | 15.3 | 123 | 56 | 55.1 | 1 | 1 | 1 | 4 | 12 | 10:50 | 11:17 |
| 67 | 19 | Alcala Pt. | 17 | 2 | 49 | 11 | 30.9 | 123 | 52 | 36.4 | 2 | 1 | 0 | 4 | 42 | 11:40 | 12:13 |
| 68 | 19 | Alcala Pt. | 17 | 1 | 49 | 13 | 1.0 | 123 | 52 | 58.7 | 1 | 4 | 1 | 4 | 13 | 12:20 | 13:00 |
| 69 | 303 | Round Rk. | 17 | 1 | 49 | 17 | 22.7 | 124 | 4 | 7.6 | 1 | 2 | 1 | 4 | 11 | 13:30 | 14:10 |
| 70 | 303 | Round Rk. | 17 | 2 | 49 | 17 | 34.4 | 124 | 4 | 33.4 | 1 | 2 | 1 | 4 | 27 | 14:20 | 14:44 |
| 71 | 304 | Entrance I. | 17 | 1 | 49 | 18 | 48.9 | 124 | 8 | 48.2 | 4 | 3 | 1 | 4 | 16 | 8:48 | 9:29 |
| 72 | 304 | Entrance I. | 17 | 2 | 49 | 18 | 42.7 | 124 | 9 | 15.3 | 4 | 3 | 1 | 4 | | 9:30 | 10:19 |
| 73 | 305 | Malaspina Pt. | 17 | 1 | 48 | 39 | 44.5 | 123 | 20 | 29.0 | 3 | 2 | 1 | 4 | 11 | 10:35 | 11:17 |
| 74 | 305 | Malaspina Pt. | 17 | 2 | 48 | 39 | 44.5 | 123 | 20 | 29.0 | 3 | 2 | 1 | 4 | 29 | 11:19 | 11:36 |
| 75 | 306 | Snake I. | 17 | 1 | 48 | 40 | 10.0 | 123 | 22 | 58.1 | 3 | 2 | 1 | 4 | 17 | 11:45 | 12:41 |
| 76 | 302 | DeCourcy I. | 17 | 1 | 48 | 40 | 10.0 | 123 | 22 | 58.1 | 2 | 1 | 1 | 1 | 7 | 11:40 | 12:17 |
| 77 | 302 | DeCourcy I. | 17 | 2 | 48 | 46 | 43.8 | 123 | 19 | 17.5 | 2 | 1 | 1 | 1 | 40 | 12:20 | 13:00 |
| 78 | 303 | Round Rk. | 17 | 2 | 48 | 46 | 43.8 | 123 | 19 | 17.5 | 2 | 1 | 1 | 1 | 37 | 14:20 | 14:50 |
| 79 | 303 | Round Rk. | 17 | 1 | 48 | 44 | 10.1 | 123 | 14 | 4.3 | 2 | 1 | 1 | 4 | 10 | 15:00 | 15:35 |
| 80 | 307 | Five Fingers | 17 | 1 | 48 | 44 | 10.1 | 123 | 14 | 4.3 | 4 | 2 | 1 | 3 | 15 | 9:20 | 10:11 |
| 81 | 307 | Five Fingers | 17 | 2 | 48 | 43 | 57.0 | 123 | 12 | 0.4 | 4 | 2 | 2 | 3 | 33 | 10:16 | 10:47 |
| 82 | 308 | Hudson Rk. | 17 | 2 | 48 | 32 | 21.3 | 123 | 17 | 13.5 | 4 | 3 | 1 | 3 | 40 | 11:27 | 12:09 |
| 83 | 308 | Hudson Rk. | 17 | 1 | 48 | 32 | 21.3 | 123 | 17 | 13.5 | 4 | 1 | 1 | 3 | 18 | 12:13 | 13:00 |
| 84 | 309 | Neck Pt. | 17 | 1 | 48 | 37 | 12.6 | 123 | 16 | 8.8 | 4 | 1 | 1 | 3 | 10 | 13:40 | 14:46 |
| 85 | 309 | Neck Pt. | 17 | 2 | 48 | 37 | 12.6 | 123 | 16 | 8.8 | 4 | 1 | 1 | 3 | 39 | 14:49 | 15:23 |
| 86 | 305 | Malaspina Pt. | 17 | 2 | 48 | 39 | 42.0 | 123 | 17 | 54.8 | 2 | 2 | 1 | 1 | 35 | 8:48 | 9:35 |
| 87 | 306 | Snake I. | 17 | 2 | 48 | 39 | 42.0 | 123 | 17 | 54.8 | 3 | 2 | 2 | 1 | 30 | 9:48 | 10:24 |
| 88 | 304 | Entrance I. | 17 | 1 | 48 | 44 | 44.6 | 123 | 24 | 32.1 | 4 | 2 | 1 | 1 | 9 | 10:40 | 11:13 |
| 89 | 310 | Grey Rk. | 17 | 1 | 48 | 45 | 1.2 | 123 | 24 | 3.0 | 2 | 1 | 1 | 1 | | 11:48 | 12:45 |
| 90 | 310 | Grey Rk. | 17 | 2 | 48 | 43 | 42.0 | 123 | 25 | 50.8 | 2 | 1 | 1 | 1 | 35 | 12:52 | 13:28 |
| 91 | 311 | Douglas I. | 17 | 2 | 48 | 43 | 42.0 | 123 | 25 | 50.8 | 2 | 1 | 1 | 1 | 43 | 13:50 | 14:30 |
| 92 | 311 | Douglas I. | 17 | 1 | 48 | 42 | 7.3 | 123 | 31 | 17.1 | 2 | 1 | 1 | 1 | 15 | 14:38 | 15:26 |

Appendix Table 2. Effort data by set (depth strata 1=0-25m; depth strata 2=26-50 m) for each fisher (F1-F5) and number of fish caught by species. Species codes are as follows: LC=lingcod, CR=copper rockfish, QR=quillback rockfish, YR=yelloweye rockfish, KG=kelp greenling, BS=buffalo sculpin, RI=red Irish lord, GS=great sculpin, CA=cabezon, PS=Pacific sanddab, SS=speckled sanddab, RS=rock sole, SD=spiny dogfish.

| Set | Depth | Effort (min) | | | | | Total | Total | Number of fish caught | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | strata | F1 | F2 | F3 | F4 | F5 | effort | fish | LC | CR | QR | YR | KG | BS | RI | GS | CA | PS | SS | RS | SD |
| 1 | 1 | 17.8 | 10.3 | 21.8 | 20.0 | | 70.0 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 2 | 2 | 17.5 | 0.0 | 20.3 | 24.0 | | 61.8 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 3 | 1 | 19.0 | 6.1 | 23.1 | 13.2 | | 61.4 | 3 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| 4 | 2 | 13.7 | 15.3 | 19.0 | 15.1 | | 63.0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| 5 | 1 | 20.7 | 0.0 | 20.4 | 20.6 | | 61.6 | 6 | 0 | 2 | 1 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 | 1 | 0 |
| 6 | 2 | 17.5 | 12.7 | 21.8 | 10.2 | | 62.2 | 15 | 0 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 11 | 0 | 1 | 0 |
| 7 | 1 | 18.4 | 15.2 | 15.2 | 11.9 | | 60.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 8 | 2 | 15.3 | 8.0 | 18.2 | 15.1 | | 56.6 | 2 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 9 | 1 | 18.5 | 10.6 | 24.5 | 11.4 | | 65.0 | 8 | 6 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 10 | 1 | 20.4 | 4.4 | 18.9 | 19.6 | | 63.3 | 9 | 4 | 1 | 0 | 0 | 2 | 0 | 2 | 0 | 0 | 0 | 0 | 0 | 0 |
| 11 | 2 | 22.6 | 18.4 | 20.8 | 0.0 | | 61.8 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 |
| 12 | 1 | 21.0 | 15.1 | 17.0 | 10.0 | | 63.1 | 10 | 9 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 13 | 2 | 21.8 | 21.2 | 9.6 | 20.1 | | 72.7 | 6 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 4 | 0 |
| 14 | 1 | 16.0 | 11.6 | 20.5 | 16.5 | | 64.5 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| 15 | 2 | 26.0 | 3.4 | 19.0 | 19.0 | | 67.5 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 16 | 1 | 11.7 | 8.4 | 23.2 | 24.9 | | 68.0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 17 | 2 | 19.5 | 10.0 | 20.1 | 15.7 | | 65.3 | 3 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 |
| 18 | 1 | 21.9 | 11.0 | 18.9 | 8.5 | | 60.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 19 | 2 | 13.6 | 0.0 | 17.1 | 21.2 | | 51.8 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 20 | 2 | 21.5 | 12.8 | 20.0 | 19.5 | | 73.8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 21 | 1 | 19.0 | 10.3 | 14.7 | 16.5 | | 60.4 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 22 | 1 | 21.9 | 7.9 | 20.0 | 11.1 | | 60.8 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 23 | 2 | 15.7 | 15.4 | 13.4 | 19.4 | | 63.8 | 4 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 24 | 1 | 16.1 | 14.9 | 14.5 | 14.7 | | 60.2 | 3 | 1 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 25 | 2 | 13.4 | 11.9 | 19.6 | 17.0 | | 61.9 | 6 | 1 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 26 | 2 | 18.5 | 13.1 | 17.1 | 14.6 | | 63.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 27 | 1 | 14.5 | 10.5 | 19.6 | 18.9 | | 63.5 | 3 | 1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 28 | 1 | 19.0 | 16.8 | 19.4 | 14.1 | | 69.2 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 29 | 2 | 17.1 | 20.3 | 20.1 | 10.0 | | 67.4 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 30 | 1 | 19.4 | 17.6 | 17.4 | 13.0 | | 67.4 | 3 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |

Appendix Table 2

| Set | Depth strata | Effort (min) F1 | F2 | F3 | F4 | F5 | Total effort | Total fish | Species LC | CR | QR | YR | KG | BS | RI | GS | CA | PS | SS | RS | SD |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 31 | 2 | 17.0 | 14.9 | 19.0 | 6.9 | | 57.8 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 32 | 2 | 20.1 | 2.5 | 19.4 | 19.1 | | 61.1 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 1 | 0 |
| 33 | 1 | 22.7 | 17.3 | 9.8 | 10.8 | | 60.5 | 6 | 0 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 35 | 2 | 0.0 | 6.3 | 8.6 | 11.2 | | 26.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 37 | 1 | 18.1 | 7.0 | 17.2 | 19.8 | | 62.1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 38 | 2 | 14.6 | 12.3 | 20.1 | 20.0 | | 67.0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 3 |
| 39 | 1 | 20.9 | 10.0 | 18.4 | 16.9 | | 66.2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 40 | 2 | 17.1 | 12.3 | 18.5 | 16.0 | | 63.9 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 |
| 41 | 2 | 25.0 | 9.0 | 18.2 | 14.5 | | 66.7 | 8 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 |
| 42 | 1 | 15.7 | 12.0 | 17.0 | 17.0 | | 61.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 43 | 1 | 17.1 | 16.0 | 16.6 | 12.4 | | 62.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 44 | 2 | 10.1 | 13.6 | 21.4 | 19.0 | | 64.1 | 4 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 45 | 1 | 16.3 | 18.8 | 13.7 | 14.3 | | 63.1 | 6 | 6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 46 | 2 | 18.0 | 19.3 | 17.5 | 6.0 | | 60.8 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 47 | 1 | 21.0 | 20.1 | 19.7 | 0.0 | | 60.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 48 | 2 | 23.4 | 12.1 | 25.4 | 0.0 | | 60.8 | 6 | 4 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 49 | 1 | 21.3 | 0.0 | 18.1 | 21.2 | | 60.6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 50 | 2 | 8.8 | 0.0 | 14.8 | 13.9 | | 37.5 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 51 | 1 | 16.1 | 23.7 | 22.6 | 4.0 | | 66.3 | 4 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 52 | 2 | 19.3 | 21.0 | 20.1 | 2.7 | | 63.0 | 4 | 1 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 53 | 1 | 15.8 | 18.9 | 17.0 | 9.2 | | 60.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 54 | 2 | 16.7 | 17.8 | 15.8 | 14.0 | | 64.3 | 3 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 55 | 1 | 12.9 | 19.7 | 19.5 | 11.6 | | 63.7 | 5 | 5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 56 | 2 | 13.7 | 15.2 | 16.6 | 16.1 | | 61.4 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 57 | 2 | 18.3 | 12.0 | 16.6 | 14.9 | | 61.8 | 11 | 6 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 3 | 0 | 1 | 0 |
| 58 | 1 | 17.7 | 10.9 | 19.2 | 12.8 | | 60.6 | 4 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 59 | 2 | 19.5 | 14.6 | 20.0 | 5.7 | | 59.8 | 2 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 60 | 1 | 20.1 | 10.5 | 17.2 | 14.0 | | 61.8 | 4 | 4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 61 | 1 | 14.0 | 14.1 | 18.2 | 18.3 | | 64.6 | 3 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 62 | 1 | 18.9 | 13.3 | 17.0 | 16.3 | | 65.4 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 63 | 2 | 14.7 | 18.7 | 16.4 | 15.2 | | 64.9 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 64 | 2 | 15.0 | 13.4 | 17.3 | 14.9 | | 60.5 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 |

Appendix Table 2

| Set | Depth strata | Effort (min) | | | | | Total effort | Total fish | Species | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | F1 | F2 | F3 | F4 | F5 | | | LC | CR | QR | YR | KG | BS | RI | GS | CA | PS | SS | RS | SD |
| 65 | 1 | 12.5 | 13.7 | 17.5 | 21.4 | | 65.1 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 66 | 1 | 18.2 | 14.0 | 16.0 | 14.7 | | 62.8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 67 | 2 | 14.7 | 17.4 | 14.9 | 14.7 | | 61.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 68 | 1 | 15.3 | 20.0 | 16.2 | 14.2 | | 65.6 | 3 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 69 | 1 | 12.1 | 17.8 | 17.7 | 15.2 | | 62.8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 70 | 2 | 14.1 | 16.0 | 17.0 | 18.7 | | 65.8 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 71 | 1 | 15.3 | 14.7 | 15.0 | 0.0 | 16.5 | 61.5 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 72 | 2 | 16.8 | 16.1 | 14.0 | 0.0 | 13.0 | 59.9 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 73 | 1 | 13.5 | 16.8 | 14.0 | 1.3 | 21.3 | 66.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 74 | 2 | 0.3 | 0.0 | 4.0 | 0.0 | 3.5 | 7.8 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 75 | 1 | 13.0 | 11.8 | 15.5 | 0.0 | 16.7 | 56.9 | 3 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 76 | 1 | 18.8 | 13.0 | 18.0 | 14.0 | | 63.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 77 | 2 | 20.3 | 0.0 | 21.5 | 19.3 | | 61.0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 78 | 2 | 20.0 | 0.0 | 20.0 | 20.0 | | 60.1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 79 | 1 | 17.9 | 7.8 | 20.1 | 16.7 | | 62.4 | 1 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 80 | 1 | 15.5 | 18.6 | 15.2 | 0.0 | 17.0 | 66.3 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 0 |
| 81 | 2 | 16.0 | 14.8 | 16.2 | 0.0 | 12.7 | 59.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 82 | 2 | 18.5 | 9.5 | 18.7 | 0.0 | 18.1 | 64.7 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 |
| 83 | 1 | 12.0 | 20.1 | 19.4 | 0.0 | 14.0 | 65.5 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| 84 | 1 | 11.3 | 15.1 | 18.0 | 0.0 | 19.3 | 63.7 | 17 | 4 | 2 | 10 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 85 | 2 | 20.0 | 16.8 | 13.0 | 0.0 | 13.8 | 63.7 | 4 | 1 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 86 | 2 | 13.0 | 14.3 | 17.6 | 0.0 | 16.5 | 61.4 | 2 | 2 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 87 | 2 | 16.5 | 12.5 | 13.9 | 0.0 | 21.0 | 63.9 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 88 | 1 | 17.5 | 18.1 | 15.0 | 0.0 | 16.0 | 66.6 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 89 | 1 | 12.0 | 16.8 | 15.8 | 0.0 | 17.0 | 61.6 | 5 | 0 | 4 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 90 | 2 | 16.0 | 13.3 | 16.0 | 0.0 | 17.0 | 62.3 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 91 | 2 | 15.7 | 13.0 | 16.0 | 0.0 | 17.0 | 61.7 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 92 | 1 | 19.0 | 18.2 | 18.0 | 0.0 | 6.8 | 62.0 | 4 | 2 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Total | | 1514 | 1141 | 1586 | 1044 | 278 | 5561.5 | 232 | 102 | 35 | 31 | 2 | 5 | 2 | 4 | 1 | 7 | 24 | 1 | 9 | 9 |

**Appendix Table 3a. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for lingcod caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m.**

| Set | Statistical Area | Depth Strata | Depth Caught (m) | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|
| 2 | 19 | 2 | 30 | | | | | 4 |
| 8 | 18 | 2 | 37 | 712 | 3780 | 2 | 3 | 3 |
| 9 | 18 | 1 | 10 | 584 | 1920 | 1 | 2 | 4 |
| 9 | 18 | 1 | 12 | 500 | 1090 | 1 | 2 | 1 |
| 9 | 18 | 1 | 16 | 576 | 1150 | 1 | 2 | 4 |
| 9 | 18 | 1 | 17 | 518 | 1260 | 1 | 2 | 3 |
| 9 | 18 | 1 | 13 | | | | 1 | 1 |
| 9 | 18 | 1 | 13 | | | | | 3 |
| 10 | 19 | 1 | 10.5 | 395 | 520 | 1 | 1 | 3 |
| 10 | 19 | 1 | 13.3 | 470 | 890 | 1 | 2 | 4 |
| 10 | 19 | 1 | 14 | 476 | 880 | 1 | 2 | 1 |
| 10 | 19 | 1 | 15.5 | 465 | 930 | 1 | 1 | 2 |
| 12 | 19 | 1 | 13 | 574 | 1240 | 1 | 2 | 3 |
| 12 | 19 | 1 | 16 | 491 | 890 | 1 | 1 | 1 |
| 12 | 19 | 1 | 20 | 456 | 780 | 1 | 1 | 2 |
| 12 | 19 | 1 | 20 | 490 | 1090 | 1 | 2 | 3 |
| 12 | 19 | 1 | 20 | 492 | 1040 | 1 | 2 | 4 |
| 12 | 19 | 1 | 23 | 500 | 1020 | 1 | 1 | 3 |
| 12 | 19 | 1 | 25 | 548 | 1490 | 1 | 2 | 3 |
| 12 | 19 | 1 | 13 | | | | | 1 |
| 12 | 19 | 1 | 22 | | | | | 3 |
| 13 | 19 | 2 | 42 | 576 | 1790 | 1 | 2 | 3 |
| 15 | 19 | 2 | 31.3 | 368 | 430 | 1 | 1 | 4 |
| 16 | 18 | 1 | 13 | 369 | 380 | 1 | 1 | 3 |
| 21 | 18 | 1 | 14 | 545 | 1610 | 1 | 3 | 4 |
| 22 | 19 | 1 | 24 | 455 | 790 | 1 | 1 | 3 |
| 22 | 19 | 1 | 16 | 630 | 2370 | 2 | 2 | 4 |
| 23 | 18 | 2 | 38 | 607 | 2120 | 1 | 2 | 2 |
| 24 | 18 | 1 | 18 | 624 | 2390 | 1 | 2 | 3 |
| 25 | 18 | 2 | 38 | 455 | 890 | 1 | 1 | 2 |
| 27 | 18 | 1 | 15 | 830 | 6450 | 1 | | 1 |

Appendix Table 3a.

| Set | Statistical Area | Depth Strata | Depth (m) Caught | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|
| 28 | 18 | 1 | 9 | 545 | 1790 | 1 | 3 | 1 |
| 29 | 18 | 2 | 30 | 675 | 3310 | 2 | 2 | 2 |
| 30 | 19 | 1 | 18 | 447 | 700 | 1 | 1 | 2 |
| 30 | 19 | 1 | 10 | 546 | 1600 | 2 | 2 | 2 |
| 31 | 19 | 2 | 26 | 602 | 2500 | 1 | 2 | 1 |
| 31 | 19 | 2 | 29 | 581 | 2250 | 2 | 2 | 2 |
| 41 | 19 | 2 | 39 | 608 | 1960 | 1 | 1 | 4 |
| 41 | 19 | 2 | 40 | 530 | 1520 | 1 | 2 | 2 |
| 41 | 19 | 2 | 40 | 498 | 1003 | 1 | 1 | 4 |
| 41 | 19 | 2 | 40 | 574 | 1680 | 2 | 1 | 1 |
| 41 | 19 | 2 | 40 | 570 | 1780 | 2 | 2 | 2 |
| 41 | 19 | 2 | 45 | 643 | 2570 | 2 | 2 | 1 |
| 44 | 17 | 2 | 31 | 760 | 4560 | 1 | 3 | 1 |
| 44 | 17 | 2 | 30 | 760 | 4230 | 2 | 3 | 1 |
| 44 | 17 | 2 | 41 | 726 | 3630 | 2 | 3 | 2 |
| 45 | 17 | 1 | 4 | 540 | 1580 | 1 | 3 | 3 |
| 45 | 17 | 1 | 4 | 630 | 2660 | 1 | 3 | 4 |
| 45 | 17 | 1 | 10 | 514 | 1160 | 1 | 2 | 2 |
| 45 | 17 | 1 | 10 | 628 | 2550 | 1 | 3 | 3 |
| 45 | 17 | 1 | 7 | 755 | 4280 | 2 | 3 | 1 |
| 45 | 17 | 1 | 10 | 605 | 2170 | 2 | 3 | 4 |
| 46 | 17 | 2 | 35 | 805 | 5190 | 2 | 3 | 2 |
| 46 | 17 | 2 | 37 | 868 | 7020 | 2 | 3 | 4 |
| 48 | 17 | 2 | 29 | 467 | 1010 | 1 | 2 | 1 |
| 48 | 17 | 2 | 37 | 473 | 870 | 1 | 2 | 1 |
| 48 | 17 | 2 | 37 | 462 | 890 | 1 | 2 | 2 |
| 48 | 17 | 2 | 39 | 856 | 7280 | 2 | 3 | 2 |
| 51 | 17 | 1 | 10 | 594 | 2140 | 1 | 3 | 3 |
| 51 | 17 | 1 | 14 | 450 | 780 | 1 | 1 | 2 |
| 52 | 17 | 2 | 49 | 640 | 2660 | 2 | 3 | 3 |
| 54 | 17 | 2 | 27 | 625 | 2500 | 1 | 2 | 2 |
| 55 | 17 | 1 | 6 | 645 | 3160 | 1 | 4 | 4 |

Appendix Table 3a.

| Set | Statistical Area | Depth Strata | Depth (m) Caught | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|
| 55 | 17 | 1 | 7 | 484 | 1120 | 1 | 2 | 2 |
| 55 | 17 | 1 | 8 | 488 | 1050 | 1 | 3 | 3 |
| 55 | 17 | 1 | 15 | 594 | 1770 | 1 | 3 | 3 |
| 55 | 17 | 1 | 16 | 680 | 2920 | 2 | 3 | 1 |
| 56 | 17 | 2 | 27 | 494 | 1000 | 1 | 1 | 2 |
| 57 | 17 | 2 | 31 | 447 | 730 | 1 | 1 | 2 |
| 57 | 17 | 2 | 31 | 520 | 1180 | 1 | 2 | 4 |
| 57 | 17 | 2 | 35 | 495 | 980 | 1 | 2 | 3 |
| 57 | 17 | 2 | 38 | 448 | 760 | 1 | 1 | 2 |
| 57 | 17 | 2 | 29 | 522 | 1190 | 2 | 1 | 4 |
| 57 | 17 | 2 | 31 | 510 | 980 | 2 | 1 | 3 |
| 58 | 17 | 1 | 17 | 743 | 4250 | 1 | 3 | 2 |
| 58 | 17 | 1 | 10 | 516 | 1140 | 2 | 1 | 3 |
| 59 | 17 | 2 | 47 | 453 | 820 | 1 | 2 | 2 |
| 60 | 17 | 1 | 6 | 645 | 2150 | 1 | 3 | 4 |
| 60 | 17 | 1 | 7 | 565 | 1140 | 1 | 3 | 2 |
| 60 | 17 | 1 | 9 | 515 | 1180 | 1 | 2 | 2 |
| 60 | 17 | 1 | 9 | 682 | 3310 | 1 | 3 | 3 |
| 61 | 17 | 1 | 12 | 670 | 3240 | 1 | 4 | 2 |
| 61 | 17 | 1 | 12 | 645 | 2620 | 2 | 1 | 1 |
| 61 | 17 | 1 | 14 | 568 | 1630 | 2 | 1 | 3 |
| 63 | 17 | 1 | 25 | 474 | 960 | 1 | 1 | 4 |
| 63 | 17 | 2 | 30 | 602 | 2020 | 2 | 2 | 1 |
| 68 | 17 | 1 | 17 | 710 | 4060 | 1 | 5 | 1 |
| 68 | 17 | 1 | 16 | 480 | 1050 | 2 | 1 | 4 |
| 70 | 17 | 2 | 27 | 435 | 630 | 1 | 2 | 2 |
| 72 | 17 | 2 | 42 | 755 |  | 1 |  | 3 |
| 74 | 17 | 2 | 30 | 446 | 720 | 2 | 1 | 1 |
| 75 | 17 | 1 | 7 | 570 |  | 1 |  | 1 |
| 75 | 17 | 1 | 10 | 560 |  | 1 |  | 3 |
| 84 | 17 | 1 | 15 | 554 | 1590 | 2 | 2 | 2 |
| 84 | 17 | 1 | 16 | 537 | 1450 | 2 | 2 | 1 |

Appendix Table 3a.

| Set | Statistical Area | Depth Strata | Depth (m) | Caught | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|---|
| 84 | 17 | 1 | 19 | | 471 | 850 | 2 | 1 | 1 |
| 85 | 17 | 2 | 34 | | 503 | 1190 | 1 | 3 | 2 |
| 84 | 17 | 1 | 21 | | 815 | 5290 | 2 | 3 | 5 |
| 86 | 17 | 2 | 27 | | 484 | 1030 | 1 | 2 | 5 |
| 86 | 17 | 2 | 28 | | 405 | 570 | 1 | 1 | 1 |
| 92 | 17 | 1 | 15 | | 535 | 1540 | 1 | 3 | 2 |
| 92 | 17 | 1 | 23 | | 430 | 700 | 1 | 2 | 1 |

**Apendix Table 3b. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for copper rockfish caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m.**

| Set | Statistical Area | Depth Strata | Depth (m) Caught | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|
| 1 | 19 | 1 | 20 | 310 | 560 | 1 | 2 | 3 |
| 5 | 18 | 1 | 24.5 | 275 | 370 | 1 | 2 | 1 |
| 5 | 18 | 1 | 10.1 | 220 | 210 | 1 | 2 | 4 |
| 6 | 18 | 2 | 34.5 | 360 | 850 | 1 | 4 | 4 |
| 10 | 19 | 1 | 13 | 311 | 580 | 1 | 4 | 1 |
| 12 | 19 | 1 | 25 | 380 | 1030 | 1 | 4 | 1 |
| 19 | 18 | 2 | 28 | 313 | 450 | 2 | 2 | 4 |
| 24 | 18 | 1 | 16.5 | 354 | 750 | 1 | 3 | 1 |
| 25 | 18 | 2 | 45 | 359 | 785 | 1 | 7 | 1 |
| 25 | 18 | 2 | 30 | 301 | 630 | 2 | 7 | 3 |
| 25 | 18 | 2 | 29 | 388 | 970 | 2 | 7 | 4 |
| 25 | 18 | 2 | 29 | 355 | 675 | 1 | 4 | 2 |
| 27 | 18 | 1 | 16 | 275 | 340 | 1 | 4 | 3 |
| 27 | 18 | 1 | 14 | 315 | 555 | 2 | 3 | 1 |
| 28 | 18 | 1 | 14 | 268 | 320 | 2 | 3 | 3 |
| 33 | 18 | 1 | 10 | 265 | 310 | 1 | 4 | 1 |
| 33 | 18 | 1 | 10 | 327 | 630 | 1 | 2 | 4 |
| 33 | 18 | 1 | 10 | 280 | 380 | 2 | 2 | 2 |
| 33 | 18 | 1 | 9 | 330 | 620 | 1 | 3 | 3 |
| 33 | 18 | 1 | 10 | 266 | 280 | 1 | 4 | 2 |
| 33 | 18 | 1 | 10 | 291 | 410 | 2 | 2 | 1 |
| 37 | 19 | 1 | 18 | 239 | 250 | 2 | 1 | 4 |
| 51 | 17 | 1 | 15 | 281 | 360 | 1 | 2 | 2 |
| 51 | 17 | 1 | 16 | 382 | 1020 | 1 | 4 | 1 |
| 58 | 17 | 1 | 12 | 314 | 590 | 1 | 4 | 3 |
| 65 | 17 | 1 | 18 | 320 | 650 | 2 | 2 | 2 |
| 68 | 17 | 1 | 13 | 290 | 400 | 1 | 3 | 4 |
| 79 | 17 | 1 | 7 | 345 | | | | 2 |
| 84 | 17 | 1 | 16 | 316 | 515 | 2 | 3 | 5 |
| 84 | 17 | 1 | 21 | 386 | | 1 | | 2 |
| 89 | 17 | 1 | 20 | 375 | 825 | 1 | 4 | 5 |
| 89 | 17 | 1 | 19 | 392 | 985 | 1 | 4 | 5 |
| 89 | 17 | 1 | 19 | 368 | 980 | 1 | 4 | 5 |
| 89 | 17 | 1 | 19 | 341 | 670 | 1 | 4 | 1 |
| 92 | 17 | 1 | 15 | 305 | 550 | 2 | 2 | 5 |

**Appendix Table 3c. Fork length (mm), weight (g), sex (1=male, 2=female) and stage of maturity for quillback rockfish caught. Stage of maturity codes: 1=immature, 2=maturing, 3=maturing, 4=mature, 5=ripe, 6=spent, 7=resting. Missing data indicate the fish was not landed. Depth strata 1=0-25 m; depth strata 2=26-50 m.**

| Set | Statistical Area | Depth Strata | Depth (m) | Caught | Length (mm) | Weight (g) | Sex | Maturity | Fisher |
|---|---|---|---|---|---|---|---|---|---|
| 5 | 18 | 1 | 24.5 | | 255 | 330 | 1 | 1 | 3 |
| 6 | 18 | 2 | 38 | | 260 | 310 | 1 | 1 | 3 |
| 6 | 18 | 2 | 27 | | 279 | 390 | 2 | 2 | 3 |
| 8 | 18 | 2 | 38 | | 235 | 280 | 1 | 1 | 3 |
| 17 | 18 | 2 | 33 | | 309 | 470 | 1 | 3 | 2 |
| 23 | 18 | 2 | 38 | | 279 | 395 | 1 | 3 | 4 |
| 25 | 18 | 2 | 45 | | 323 | 740 | 1 | 3 | 4 |
| 30 | 19 | 1 | 13 | | 309 | 530 | 2 | 2 | 3 |
| 48 | 17 | 2 | 29 | | 312 | 580 | 2 | 2 | 3 |
| 52 | 17 | 1 | 25 | | 263 | 360 | 2 | 2 | 1 |
| 52 | 17 | 1 | 25 | | 363 | 880 | 1 | 2 | 2 |
| 52 | 17 | 2 | 41 | | 390 | 1110 | 2 | 2 | 4 |
| 54 | 17 | 2 | 33 | | 335 | 710 | 2 | 1 | 1 |
| 57 | 17 | 2 | 28 | | 245 | 270 | 2 | 1 | 1 |
| 58 | 17 | 1 | 10 | | 333 | 650 | 2 | 5 | 1 |
| 75 | 17 | 1 | 14 | | 360 | 890 | 2 | 5 | 5 |
| 84 | 17 | 1 | 16 | | 325 | | | | 1 |
| 84 | 17 | 1 | 17 | | 240 | 255 | 2 | 2 | 2 |
| 84 | 17 | 1 | 21 | | 272 | 360 | 1 | 3 | 2 |
| 84 | 17 | 1 | 21 | | 285 | 405 | 2 | 3 | 3 |
| 84 | 17 | 1 | 20 | | 249 | 270 | 1 | 2 | 3 |
| 84 | 17 | 1 | 20 | | 328 | 655 | 2 | 7 | 1 |
| 84 | 17 | 1 | 18 | | 321 | | 1 | | 5 |
| 84 | 17 | 1 | 21 | | 524 | | | | 1 |
| 84 | 17 | 1 | 21 | | 348 | | 2 | | 2 |
| 84 | 17 | 1 | 24 | | 348 | | 1 | | 1 |
| 85 | 17 | 2 | 34 | | 398 | 1255 | 2 | 3 | 5 |
| 85 | 17 | 2 | 34 | | 395 | 1245 | 2 | 3 | 3 |
| 85 | 17 | 2 | 42 | | 313 | 485 | 1 | 3 | 5 |
| 89 | 17 | 1 | 19 | | 257 | 320 | 1 | 2 | 1 |
| 92 | 17 | 1 | 12 | | 243 | 260 | 2 | 2 | 3 |

**Appendix Table 4. Description of maturity stages for male and female lingcod.**

| Maturity Stage | Male | Female |
|---|---|---|
| STAGE 1: Immature | • Testes are threadlike to ribbonlike.<br>• Colour is transparent white to white | • Ovaries are small and translucent.<br>• Colour is pink or white-pink.<br>• Eggs are not visible |
| STAGE 2: Maturing - small | • Testes are larger ribbonlike.<br>• Colour is white to very light brown | • Ovaries fill about 1/4 to 1/3 of body cavity.<br>• Colour is orange and opaque or semi-translucent.<br>• Blood vessels are pronounced on the ovary |
| STAGE 3: Maturing - large | • Testes fill 1/3 of body cavity.<br>• Colour is whiter than in Stage 2 | • Ovaries fill about 1/3 to 2/3 of body cavity<br>• Colour is orange.<br>• Blood vessels are pronounced on the ovary.<br>• Eggs are opaque |
| STAGE 4: Mature | • Testes fill 1/3 to 2/3 of body cavity<br>• Colour is white.<br>• A cross sectioning of testis will produce sperm at the<br>• centre of the tissue | • Ovaries fill 2/3 to 4/5 of the body cavity.<br>• Eggs are opaque.<br>• Colour is orange to white |
| STAGE 5: Ripe | • Testes fill 2/3 or more of the body cavity.<br>• Colour is white.<br>• Testis lobes are fully developed.<br>• Sperm is released from the vent with slight pressure<br>• on the body exterior. | • Ovaries fill near all of the body cavity.<br>• Colour is opaque orange to white.<br>• Eggs may be loose inside the ovary |
| STAGE 6: Spent | • Testes are moderate in size<br>• Colour is tan brown with some white colour still<br>• evident.<br>• A cross sectioning of testis will reveal some<br>• remaining sperm in centre of the gonad. | • Ovaries fill 1/3 to 2/3 of the body cavity.<br>• Colour is purple and may be bloodshot.<br>• Ovaries are flaccid and some eggs may remain<br>• within.<br>• Reabsorbing<br>• Recovering |
| STAGE 7: Resting | • Testes are relatively smaller and firm.<br>• Colour is tan brown | • Ovaries fill less than 1/3 of the body cavity.<br>• Colour is often pink.<br>• Ovaries are firm. |

Appendix Table 5. Codes used for sea state, tide, current and weather.

| Code | |
|---|---|
| Sea State | |
| 1 | calm |
| 2 | ripple |
| 3 | chop |
| 4 | swell |
| Tide | |
| 1 | ebb |
| 2 | flood |
| 3 | high |
| 4 | low |
| Current | |
| 0 | none |
| 1 | minimal |
| 2 | moderate |
| 3 | moderate-strong |
| Weather | |
| 1 | sun |
| 2 | rain |
| 3 | partly cloudy |
| 4 | overcast |
| 5 | high cloud |